


ASC Report No. 03/2012

Adaptive FEM with inhomogeneous dirichlet data: Convergence and quasi-optimality in \mathbb{R}^d

M. Aurada, M. Feischl, J. Kemetmüller, M. Page, D. Praetorius



Institute for Analysis and Scientific Computing
Vienna University of Technology — TU Wien
www.asc.tuwien.ac.at ISBN 978-3-902627-05-6

Most recent ASC Reports

- 02/2012 *P. Goldenits, G. Hrkac, D. Praetorius, D. Suess*
An Effective Integrator for the Landau-Lifshitz-Gilbert Equation
- 01/2012 *A. Arnold, L. Neumann, W. Hochhauser*
Stability of glued and embedded Glass Panes: Dunkerley straight Line as a conservative Estimate of superimposed buckling Coefficients
- 43/2011 *G. Hastermann, P. Lima, L. Morgado, E. Weinmüller*
Numerical Solution of the Density Profile Equation with p-Laplacians
- 42/2011 *M. Aurada, J.M. Melenk, D. Praetorius*
Mixed Conforming elements for the large-body limit in micromagnetics: a finite element approach
- 41/2011 *P. Amodio, T. Levitina, G. Settanni, E.B. Weinmüller*
On the Calculation of the Finite Hankel Transform Eigenfunctions
- 40/2011 *D.P. Hewett, S. Langdon, J.M. Melenk*
A high frequency hp boundary element method for scattering by convex polygons
- 39/2011 *A. Jüngel*
Semiconductor Device Problems
- 38/2011 *L. Neumann, A. Arnold, W. Hochhauser*
Zur Stabilität von geklebten und geklotzten Glasscheiben: Beurteilung der Dunkerley'schen Geraden zur Beulwertbestimmung
- 37/2011 *A. Dick, O. Koch, R. März, E. Weinmüller*
Collocation Schemes for Nonlinear Index 1 DAEs with a Singular Point
- 36/2011 *J.A. Carrillo, S. Hittmeir, A. Jüngel*
Cross diffusion and nonlinear diffusion preventing blow up in the Keller-Segel model

Institute for Analysis and Scientific Computing
Vienna University of Technology
Wiedner Hauptstraße 8–10
1040 Wien, Austria

E-Mail: admin@asc.tuwien.ac.at
WWW: <http://www.asc.tuwien.ac.at>
FAX: +43-1-58801-10196

ISBN 978-3-902627-05-6

© Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors.



ADAPTIVE FEM WITH INHOMOGENEOUS DIRICHLET DATA: CONVERGENCE AND QUASI-OPTIMALITY IN \mathbb{R}^d

M. AURADA, M. FEISCHL, J. KEMETMÜLLER, M. PAGE, AND D. PRAETORIUS

ABSTRACT. We consider the solution of second order elliptic PDEs in \mathbb{R}^d with inhomogeneous Dirichlet data by means of an h -adaptive FEM with fixed polynomial order $p \in \mathbb{N}$. As model example serves the Poisson equation with mixed Dirichlet-Neumann boundary conditions, where the inhomogeneous Dirichlet data are discretized by use of a stable projection, for instance, the L^2 -projection for $p = 1$ or the Scott-Zhang projection for general $p \geq 1$. For error estimation, we use a residual error estimator which includes the Dirichlet data oscillations. We prove convergence of the adaptive algorithm even with quasi-optimal convergence rate. Numerical experiments conclude the work.

1. INTRODUCTION

Recently, there has been a major breakthrough in the thorough mathematical understanding of convergence and quasi-optimality of h -adaptive FEM for second-order elliptic PDEs. However, the focus of the numerical analysis usually lies on model problems with homogeneous Dirichlet conditions, i.e. $\Delta u = f$ in Ω with $u = 0$ on $\Gamma = \partial\Omega$, see e.g. [CKNS, D, KS, MNS, S07]. Instead, our model problem

$$\begin{aligned} -\Delta u &= f && \text{in } \Omega, \\ u &= g && \text{on } \Gamma_D, \\ \partial_n u &= \phi && \text{on } \Gamma_N \end{aligned} \tag{1}$$

considers inhomogeneous mixed Dirichlet-Neumann boundary conditions. Here, Ω is a bounded Lipschitz domain in \mathbb{R}^d with polyhedral boundary $\Gamma = \partial\Omega$ which is split into two relatively open boundary parts, namely the Dirichlet boundary Γ_D and the Neumann boundary Γ_N , i.e. $\Gamma_D \cap \Gamma_N = \emptyset$ and $\overline{\Gamma_D} \cup \overline{\Gamma_N} = \Gamma$. We stress that the surface measure of the Dirichlet boundary has to be positive $|\Gamma_D| > 0$, whereas Γ_N is allowed to be empty. The given data formally satisfy $f \in \tilde{H}^{-1}(\Omega)$, $g \in H^{1/2}(\Gamma_D)$, and $\phi \in H^{-1/2}(\Gamma_N)$. As is usually required to derive (localized) a posteriori error estimators, we assume additional regularity of the given data, namely $f \in L^2(\Omega)$, $g \in H^1(\Gamma_D)$, and $\phi \in L^2(\Gamma_N)$.

We stress that —using results available in the literature— it is easily possible to generalize the analysis from the Laplacian $L = -\Delta$ to general symmetric and uniformly elliptic differential operators of second order. The reader is referred to the seminal work [CKNS] which treats the case of homogeneous Dirichlet data $g = 0$ and provides the necessary tools to cover general L .

Date: January 20, 2012.

2000 Mathematics Subject Classification. 65N30, 65N50.

Key words and phrases. adaptive finite element method, convergence analysis, quasi-optimality, inhomogeneous Dirichlet data.

Whereas certain work on a posteriori error estimation for (1) has been done, cf. [BCD, SV], none of the proposed adaptive algorithms have been proven to converge. While the inclusion of inhomogeneous Neumann conditions ϕ into the convergence analysis seems to be obvious, incorporating inhomogeneous Dirichlet conditions g is technically more demanding. First, this is due to the fractional-order Sobolev space $H^{1/2}(\Gamma_D)$. However, we may use techniques from adaptive boundary element methods [CMS, FKMP, KOP] to localize the non-local $H^{1/2}$ -norm in terms of a locally weighted H^1 -seminorm. Second, the given Dirichlet data have to be discretized appropriately. In an earlier work [FPP], we considered lowest-order finite elements $p = 1$ in 2D and nodal interpolation to discretize g . However, this situation is very special in the sense that our entire analysis in [FPP] is strictly bound to the lowest-order case and cannot be generalized to \mathbb{R}^d , since nodal interpolation is well-defined if and only if $d = 2$. In this work, we consider finite elements of piecewise polynomial order $p \geq 1$ and dimension $d \geq 2$. We show that each uniformly $H^{1/2}(\Gamma_D)$ -stable projection \mathbb{P}_ℓ onto the discrete trace space will do the job. However, and in contrast to the case $g = 0$, the discrete ansatz spaces \mathbb{V}_ℓ are non-nested, i.e. $\mathbb{V}_\ell \not\subseteq \mathbb{V}_{\ell+1}$. We therefore lose the orthogonality in energy norm which seems to be a major problem. The remedy is to concentrate on a quasi-Pythagoras theorem and a stronger marking criterion. The latter implies (quasi-local) equivalence of error estimators for different discretizations of the Dirichlet data. To obtain contraction of our AFEM, we may then consider (theoretically) the $H^{1/2}(\Gamma_D)$ -orthogonal projection. To obtain optimality of the marking strategy, we may consider the Scott-Zhang projection instead. Both auxiliary problems are somehow sufficiently close to the original problem with projection \mathbb{P}_ℓ , which is enforced by the marking strategy.


2. ADAPTIVE ALGORITHM

It is well-known that the Poisson problem (1) admits a unique weak solution $u \in H^1(\Omega)$ with $u = g$ on Γ_D in the sense of traces which solves the variational formulation

$$\langle \nabla u, \nabla v \rangle_\Omega = \langle f, v \rangle_\Omega + \langle \phi, v \rangle_{\Gamma_N} \quad \text{for all } v \in H_D^1(\Omega). \quad (2)$$

Here, the test space reads $H_D^1(\Omega) = \{v \in H^1(\Omega) : v = 0 \text{ on } \Gamma_D \text{ in the sense of traces}\}$, and $\langle \cdot, \cdot \rangle$ denotes the respective L^2 -scalar products. The proof relies essentially on a reformulation of (1) as a problem with homogeneous Dirichlet data via a so-called lifting operator \mathcal{L} , i.e. $\mathcal{L} : H^{1/2}(\Gamma) \rightarrow H^1(\Omega)$ is a linear and continuous operator with $(\mathcal{L}\hat{g})|_\Gamma = \hat{g}$ for all $\hat{g} \in H^{1/2}(\Gamma)$ in the sense of traces. However, although \mathcal{L} is constructed analytically, it is hardly accessible numerically in general and thus this approach is not feasible in practice.

This section provides an overview on this work and its main results. We analyze a common adaptive mesh-refining algorithm of the type



which is stated in detail below in Section 2.5. We start with a discussion of its four modules.

2.1. The module solve. Let \mathcal{T}_ℓ be a regular triangulation of Ω into simplices, i.e. tetrahedra for 3D resp. triangles for 2D, which is generated from an initial triangulation \mathcal{T}_0 . Let \mathcal{E}_ℓ be the set of facets, i.e. faces for 3D and edges for 2D, respectively. This set is split into interior facets $\mathcal{E}_\ell^\Omega = \{E \in \mathcal{E}_\ell : E \cap \Omega \neq \emptyset\}$, i.e. each $E \in \mathcal{E}_\ell^\Omega$ satisfies $E = T_+ \cap T_-$ for $T_\pm \in \mathcal{T}_\ell$, as well as boundary facets $\mathcal{E}_\ell^\Gamma = \mathcal{E}_\ell \setminus \mathcal{E}_\ell^\Omega$. We assume that the partition of Γ into Dirichlet boundary Γ_D and Neumann boundary Γ_N is resolved, i.e. \mathcal{E}_ℓ^Γ is split into

$\mathcal{E}_\ell^D = \{E \in \mathcal{E}_\ell : E \subseteq \bar{\Gamma}_D\}$ and $\mathcal{E}_\ell^N = \{E \in \mathcal{E}_\ell : E \subseteq \bar{\Gamma}_N\}$. Note that \mathcal{E}_ℓ^D (resp. \mathcal{E}_ℓ^N) provides a regular triangulation of the $(d-1)$ -dimensional manifold Γ_D (resp. Γ_N).

We use conforming elements of fixed polynomial order $p \in \mathbb{N}$, where the ansatz space reads

$$\mathcal{S}^p(\mathcal{T}_\ell) = \{V_\ell \in C(\bar{\Omega}) : V_\ell|_T \text{ is a polynomial of degree at most } p \text{ on } T \in \mathcal{T}_\ell\}. \quad (3)$$

Since a discrete function $U_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ cannot satisfy general continuous Dirichlet conditions, we have to discretize the given data $g \in H^1(\Gamma_D)$. To this purpose, let $\mathbb{P}_\ell : H^{1/2}(\Gamma_D) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell^D)$ be a projection onto the discrete trace space

$$\mathcal{S}^p(\mathcal{E}_\ell^D) = \{V_\ell|_{\Gamma_D} : V_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)\} \quad (4)$$

As in the continuous case, it is well-known that there is a unique $U_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ with $U_\ell = \mathbb{P}_\ell g$ on Γ_D which solves the Galerkin formulation

$$\langle \nabla U_\ell, \nabla V_\ell \rangle_\Omega = \langle f, V_\ell \rangle_\Omega + \langle \phi, V_\ell \rangle_{\Gamma_N} \quad \text{for all } V_\ell \in \mathcal{S}_D^p(\mathcal{T}_\ell). \quad (5)$$

Here, the test space is given by $\mathcal{S}_D^p(\mathcal{T}_\ell) = \mathcal{S}^p(\mathcal{T}_\ell) \cap H_D^1(\Omega) = \{V_\ell \in \mathcal{S}^p(\mathcal{T}_\ell) : V_\ell = 0 \text{ on } \Gamma_D\}$. We assume that `solve` computes the exact Galerkin solution of (5). Arguing as e.g. in [BM, S07], it is, however, possible to include an approximate solver into our analysis.

Possible choices for \mathbb{P}_ℓ include the L^2 -orthogonal projection for the lowest-order case $p = 1$, which is considered in [BCD], or the Scott-Zhang projection from [SZ] which is proposed in [SV]. Our analysis below will also theoretically use the $H^{1/2}(\Gamma_D)$ -orthogonal projection.

2.2. The module estimate. We start with the element data oscillations

$$\text{osc}_{\mathcal{T},\ell}^2 := \sum_{T \in \mathcal{T}_\ell} \text{osc}_{\mathcal{T},\ell}(T)^2, \text{ where } \text{osc}_{\mathcal{T},\ell}(T)^2 := |T|^{2/d} \|(1 - \Pi_T)(f + \Delta U_\ell)\|_{L^2(T)}^2 \quad (6)$$

and where $\Pi_T : L^2(T) \rightarrow \mathcal{P}^{p-1}(T)$ denotes the L^2 -orthogonal projection. These arise in the efficiency estimate for residual error estimators. Moreover, the efficiency involves the Neumann data oscillations

$$\text{osc}_{N,\ell}^2 := \sum_{E \in \mathcal{E}_\ell^N} \text{osc}_{N,\ell}(E)^2, \text{ where } \text{osc}_{N,\ell}(E)^2 := |T|^{1/d} \|(1 - \Pi_E)\phi\|_{L^2(E)}^2 \quad (7)$$

with $T \in \mathcal{T}_\ell$ being the unique element with $E \subseteq \partial T$ and where $\Pi_E : L^2(E) \rightarrow \mathcal{P}^{p-1}(E)$ denotes the L^2 -orthogonal projection on the boundary. Finally, the approximation of the Dirichlet data $\mathbb{P}_\ell g \approx g \in H^1(\Gamma_D)$ is controlled by the Dirichlet data oscillations

$$\text{osc}_{D,\ell}^2 := \sum_{E \in \mathcal{E}_\ell^D} \text{osc}_{D,\ell}(E)^2, \text{ where } \text{osc}_{D,\ell}(E)^2 := |T|^{1/d} \|(1 - \Pi_E)\nabla_\Gamma g\|_{L^2(E)}^2, \quad (8)$$

where again $T \in \mathcal{T}_\ell$ denotes the unique element with $E \subseteq \partial T$. Moreover, $\nabla_\Gamma(\cdot)$ denotes the surface gradient. We recall that up to shape regularity we have equivalence $|T|^{1/d} \simeq \text{diam}(T)$ as well as $\text{diam}(T) \simeq \text{diam}(E)$ for all $T \in \mathcal{T}_\ell$ and $E \in \mathcal{E}_\ell$ with $E \subseteq \partial T$.

We use a residual error estimator $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ which is split into general contributions and Dirichlet oscillations, i.e.

$$\varrho_\ell^2 = \sum_{T \in \mathcal{T}_\ell} \varrho_\ell(T)^2 \quad (9)$$

with corresponding refinement indicators

$$\varrho_\ell(T)^2 := |T|^{2/d} \|f + \Delta U_\ell\|_{L^2(T)}^2 + |T|^{1/d} (\|[\partial_n U_\ell]\|_{L^2(\partial T \cap \Omega)}^2 + \|\phi - \partial_n U_\ell\|_{L^2(\partial T \cap \Gamma_N)}^2). \quad (10)$$

The module **estimate** returns the elementwise contributions $\varrho_\ell(T)^2$ and $\text{osc}_{D,\ell}(E)^2$ for all $T \in \mathcal{T}_\ell$ and $E \in \mathcal{E}_\ell^D$.

2.3. The module mark. For element marking, we use a modification of the Dörfler marking [D] proposed firstly in Stevenson [S07]. In each step of the adaptive loop, we mark either elements or Dirichlet edges for refinement, where the latter is only done if $\text{osc}_{D,\ell}$ is large when compared to ϱ_ℓ . A precise statement of the module **mark** is part of Algorithm 1 below.

2.4. The module refine. Locally refined meshes are obtained by use of the newest vertex bisection algorithm, see e.g. [S08, T], where $\mathcal{T}_{\ell+1} = \text{refine}(\mathcal{T}_\ell, \mathcal{M}_\ell)$ for a set $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell$ of marked elements returns the coarsest regular triangulation $\mathcal{T}_{\ell+1}$ such that all marked elements $T \in \mathcal{M}_\ell$ have been refined by at least one bisection. Arguing as in [KS], one may also use variants of newest vertex bisection, where each $T \in \mathcal{M}_\ell$ is refined by at least n bisections with arbitrary, but fixed $n \in \mathbb{N}$.

2.5. Adaptive loop. With the foregoing modules, the adaptive mesh-refining algorithm takes the following form.

Algorithm 1. *Let adaptivity parameters $0 < \theta_1, \theta_2, \vartheta < 1$ and initial triangulation \mathcal{T}_0 be given. For each $\ell = 0, 1, 2, \dots$ do:*

- (i) *Compute discrete solution $U_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$.*
- (ii) *Compute refinement indicators $\varrho_\ell(T)$ and $\text{osc}_{D,\ell}(E)$ for all $T \in \mathcal{T}_\ell$ and $E \in \mathcal{E}_\ell^D$.*
- (iii) *Provided that $\text{osc}_{D,\ell}^2 \leq \vartheta \varrho_\ell^2$, choose $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell$ such that*

$$\theta_1 \varrho_\ell^2 \leq \sum_{T \in \mathcal{M}_\ell} \varrho_\ell(T)^2. \quad (11)$$

- (iv) *Provided that $\text{osc}_{D,\ell}^2 > \vartheta \varrho_\ell^2$, choose $\mathcal{M}_\ell^D \subseteq \mathcal{E}_\ell^D$ such that*

$$\theta_2 \text{osc}_{D,\ell}^2 \leq \sum_{E \in \mathcal{M}_\ell^D} \text{osc}_{D,\ell}(E)^2 \quad (12)$$

and let $\mathcal{M}_\ell := \{T \in \mathcal{T}_\ell : \exists E \in \mathcal{M}_\ell^D \quad E \subseteq \partial T\}$.

- (v) *Use newest vertex bisection to generate $\mathcal{T}_{\ell+1} = \text{refine}(\mathcal{T}_\ell, \mathcal{M}_\ell)$.*
- (vi) *Update counter $\ell \mapsto \ell + 1$ and go to (i).*

2.6. Main results. Throughout, we assume that the projections $\mathbb{P}_\ell : H^{1/2}(\Gamma_D) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell^D)$ are uniformly $H^{1/2}(\Gamma_D)$ -stable, i.e. the operator norm is uniformly bounded

$$\|\mathbb{P}_\ell : H^{1/2}(\Gamma_D) \rightarrow H^{1/2}(\Gamma_D)\| \leq C_{\text{stab}} < \infty \quad (13)$$

with some ℓ -independent constant $C_{\text{stab}} > 0$. This assumption is guaranteed for the $H^{1/2}(\Gamma_D)$ -orthogonal projection with $C_{\text{stab}} = 1$. Moreover, the $L^2(\Gamma_D)$ -orthogonal projection for the lowest-order case $p = 1$ and newest vertex bisection is uniformly bounded [KPP], and so is the Scott-Zhang projection [SZ] onto $\mathcal{S}^p(\mathcal{E}_\ell^D)$ for arbitrary $p \geq 1$.

First, our discretization is quasi-optimal in the sense of the Céa lemma. Note that estimate (14) does not depend on the precise choice of \mathbb{P}_ℓ , and the minimum is taken over all discrete functions. Unlike our observation, the result in e.g. [BCD, Theorem 6.1] takes the minimum with respect to the affine space $\{W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell) : W_\ell|_{\Gamma_D} = \mathbb{P}_\ell g\}$ and for first-order $p = 1$ only.

Proposition 2 (Céa-type estimate in H^1 -norm). *The Galerkin solution satisfies*

$$\|u - U_\ell\|_{H^1(\Omega)} \leq C_{\text{Céa}} \min_{W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)} \|u - W_\ell\|_{H^1(\Omega)}. \quad (14)$$

The constant $C_{\text{Céa}} > 0$ depends only on Ω , Γ_D , shape regularity of \mathcal{T}_ℓ , the polynomial degree $p \geq 1$, and the constant $C_{\text{stab}} > 0$.

Second, the considered error estimator provides an upper bound and, up to data oscillations, also a lower bound for the Galerkin error.

Proposition 3 (reliability and efficiency of η_ℓ). *The error estimator $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ is reliable*

$$\|u - U_\ell\|_{H^1(\Omega)}^2 \leq C_{\text{rel}} \eta_\ell^2 \quad (15)$$

and efficient

$$C_{\text{eff}}^{-1} \eta_\ell^2 \leq \|\nabla(u - U_\ell)\|_{L^2(\Omega)}^2 + \text{osc}_{\mathcal{T},\ell}^2 + \text{osc}_{N,\ell}^2 + \text{osc}_{D,\ell}^2. \quad (16)$$

The constants $C_{\text{rel}}, C_{\text{eff}} > 0$ depend on Ω and Γ_D , on the polynomial degree $p \geq 1$, stability $C_{\text{stab}} > 0$, the initial triangulation \mathcal{T}_0 , and on the use of newest vertex bisection.

Note that convergence of Algorithm 1 in the sense of $\lim_\ell U_\ell = u$ in $H^1(\Omega)$ is a priori unclear since adaptive mesh-refinement does not guarantee that the local mesh-size tends uniformly to zero. However, we have the following convergence result which is proved in the frame of the estimator reduction concept from [AFP].

Theorem 4 (convergence of AFEM). (i) *Suppose that the discretization of the Dirichlet data guarantees some a priori convergence*

$$\lim_{\ell \rightarrow \infty} \|g_\infty - \mathbb{P}_\ell g\|_{H^{1/2}(\Gamma_D)} = 0 \quad (17)$$

with a certain limit $g_\infty \in H^{1/2}(\Gamma_D)$. Then, for any choice of the adaptivity parameters $0 < \theta_1, \theta_2, \vartheta < 1$, Algorithm 1 guarantees convergence

$$\lim_{\ell \rightarrow \infty} \|u - U_\ell\|_{H^1(\Omega)} = 0 \quad (18)$$

and, in particular, $g_\infty = g$.

(ii) *Assumption (17) is satisfied for the $H^{1/2}(\Gamma_D)$ -orthogonal projection, the $L^2(\Gamma_D)$ -projection for $p = 1$, and the Scott-Zhang projection for arbitrary $p \geq 1$.*

Current quasi-optimality results on AFEM rely on the fact that the estimator $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ is equivalent to some linear convergent quasi-error quantity Δ_ℓ . Whereas, the convergence theorem (Theorem 4) also holds for the usual Dörfler marking, our contraction theorem relies on Stevenson's modification (11)–(12). Moreover, we stress that the convergence theorem is constrained by the a priori convergence assumption (17), whereas the following contraction result is not.

Theorem 5 (contraction of AFEM). *Let $P_\ell : H^{1/2}(\Gamma_D) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell^D)$ be the $H^{1/2}(\Gamma_D)$ -orthogonal projection. Let $\tilde{U}_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ the Galerkin solution of (5) with $\tilde{U}_\ell|_{\Gamma_D} = P_\ell g$ and $\tilde{\eta}_\ell^2 = \tilde{\varrho}_\ell^2 + \text{osc}_{D,\ell}^2$ be the associated error estimator from (9) with U_ℓ replaced by \tilde{U}_ℓ . Then, for arbitrary $0 < \theta_1, \theta_2 < 1$ and sufficiently small $0 < \vartheta < 1$, Algorithm 1 guarantees the existence of constants $\lambda, \mu > 0$ and $0 < \kappa < 1$ such that the combined error quantity*

$$\Delta_\ell := \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + \lambda \|g - P_\ell g\|_{H^{1/2}(\Gamma_D)}^2 + \mu \tilde{\eta}_\ell^2 \geq 0 \quad (19)$$

satisfies a contraction property

$$\Delta_{\ell+1} \leq \kappa \Delta_\ell \quad \text{for all } \ell \in \mathbb{N}_0. \quad (20)$$

Moreover, there are constants $C_{\text{low}}, C_{\text{high}} > 0$ such that

$$C_{\text{low}} \Delta_\ell \leq \eta_\ell^2 \leq C_{\text{high}} \Delta_\ell. \quad (21)$$

In particular, this implies convergence $\lim_\ell \|u - U_\ell\|_{H^1(\Omega)} = 0 = \lim_\ell \eta_\ell$ of Algorithm 1 independently of the precise choice of the uniformly $H^{1/2}(\Gamma_D)$ -stable projection \mathbb{P}_ℓ .

To state our quasi-optimality result for Algorithm 1, we need to introduce further notation. Recall that, for a given triangulation \mathcal{T}_ℓ and $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell$,

$$\mathcal{T}_{\ell+1} = \text{refine}(\mathcal{T}_\ell, \mathcal{M}_\ell) \quad (22)$$

denotes the coarsest regular triangulation such that all marked elements $T \in \mathcal{M}_\ell$ have been refined by (at least one) bisection. Moreover, we write

$$\mathcal{T}_\star = \text{refine}(\mathcal{T}_\ell) \quad (23)$$

if \mathcal{T}_\star is a finite refinement of \mathcal{T}_ℓ , i.e., there are finitely many triangulations $\mathcal{T}_{\ell+1}, \dots, \mathcal{T}_n$ and sets of marked elements $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell, \dots, \mathcal{M}_{n-1} \subseteq \mathcal{T}_{n-1}$ such that $\mathcal{T}_\star = \mathcal{T}_n$ and $\mathcal{T}_{j+1} = \text{refine}(\mathcal{T}_j, \mathcal{M}_j)$ for all $j = \ell, \dots, n-1$. Finally, for a fixed initial mesh \mathcal{T}_0 , let $\mathbb{T} = \{\mathcal{T}_\star : \mathcal{T}_\star = \text{refine}(\mathcal{T}_0)\}$ be the set of all meshes which can be obtained by newest vertex bisection as well as the set $\mathbb{T}_N = \{\mathcal{T}_\star \in \mathbb{T} : \#\mathcal{T}_\star - \#\mathcal{T}_0 \leq N\}$ of all triangulations which have at most N more elements than the initial mesh \mathcal{T}_0 .

Recall that Algorithm 1 only sees the error estimator $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$, but not the error $\|u - U_\ell\|_{H^1(\Omega)}$. From this point of view, it is natural to ask for the best possible convergence rate for the error estimator. This can be characterized by means of an artificial approximation class \mathbb{A}_s : For $s \geq 0$, we write

$$(u, f, g, \phi) \in \mathbb{A}_s \quad \stackrel{\text{def}}{\iff} \quad \sup_{N \in \mathbb{N}} \inf_{\mathcal{T}_\star \in \mathbb{T}_N} N^s \eta_\star < \infty, \quad (24)$$

where $\eta_\star^2 = \varrho_\star^2 + \text{osc}_{D,\star}^2$ denotes the error estimator for the optimal mesh $\mathcal{T}_\star \in \mathbb{T}_N$. By definition, this implies that a convergence rate $\eta_\star = \mathcal{O}(N^{-s})$ is possible if the optimal meshes are chosen. The following theorem states that Algorithm 1, in fact, guarantees $\eta_\ell = \mathcal{O}(N^{-s})$ for the adaptively generated meshes \mathcal{T}_ℓ .

Theorem 6 (quasi-optimality of AFEM). *Suppose that the sets \mathcal{M}_ℓ resp. \mathcal{M}_ℓ^D in step (iii)–(iv) of Algorithm 1 are chosen with minimal cardinality. Then, for sufficiently small $0 < \theta_1, \vartheta < 1$, but arbitrary $0 < \theta_2 < 1$, Algorithm 1 guarantees the existence of a constant $C_{\text{opt}} > 0$ such that*

$$(u, f, g, \phi) \in \mathbb{A}_s \quad \iff \quad \forall \ell \in \mathbb{N} \quad \eta_\ell \leq C_{\text{opt}} (\#\mathcal{T}_\ell - \#\mathcal{T}_0)^{-s}, \quad (25)$$

i.e. each possible convergence rate $s > 0$ is, in fact, asymptotically obtained by AFEM.

We stress that, up to now and as far as the error estimator is concerned, only reliability (15) is needed for the analysis. Finally, the lower bound (16) for the error estimator allows to characterize the approximation class \mathbb{A}_s in terms of the regularity of the sought solution and the given data.

Theorem 7 (characterization of \mathbb{A}_s). *It holds $(u, f, g, \phi) \in \mathbb{A}_s$ if and only if the following four conditions hold:*

$$\sup_{N \in \mathbb{N}} \inf_{\mathcal{T}_\star \in \mathbb{T}_N} \min_{V_\star \in \mathcal{S}^p(\mathcal{T}_\star)} N^s \|u - V_\star\|_{H^1(\Omega)} < \infty, \quad (26)$$

$$\sup_{N \in \mathbb{N}} \inf_{\mathcal{T}_\star \in \mathbb{T}_N} N^s \text{osc}_{\mathcal{T}, \star} < \infty, \quad (27)$$

$$\sup_{N \in \mathbb{N}} \inf_{\mathcal{T}_\star \in \mathbb{T}_N} N^s \text{osc}_{N, \star} < \infty, \quad (28)$$

$$\sup_{N \in \mathbb{N}} \inf_{\mathcal{T}_\star \in \mathbb{T}_N} N^s \text{osc}_{D, \star} < \infty, \quad (29)$$

i.e. the estimator —and according to reliability hence the Galerkin error— converges with the best possible rate allowed by the regularity of the sought solution and the given data.

2.7. Outline. Since our analysis is strongly built on properties of the Scott-Zhang projection, Section 3 collects the essential properties of the latter. This knowledge is used to prove Proposition 2. Moreover, we prove that the Scott-Zhang error in a weighted $H^1(\Gamma_D)$ -seminorm is equivalent to the Dirichlet oscillations. This allows to prove Proposition 3 with an estimator η_ℓ which does not explicitly contain the chosen projection \mathbb{P}_ℓ . Section 4 is concerned with the proof of Theorem 4. Section 5 gives the proof for the contraction result of Theorem 5. Finally, the proof of the quasi-optimality results of Theorem 6 and Theorem 7 are found in Section 6. Some numerical experiments in Section 7 conclude the work.

In all statements, the constants involved and their dependencies are explicitly stated. In proofs, however, we use the symbol \lesssim to abbreviate \leq up to a multiplicative constant. Moreover, \simeq abbreviates that both estimates \lesssim and \gtrsim hold.

3. SCOTT-ZHANG PROJECTION

The main tool of our analysis is the Scott-Zhang projection

$$\mathbb{J}_\ell : H^1(\Omega) \rightarrow \mathcal{S}^p(\mathcal{T}_\ell) \quad (30)$$

from [SZ]. A first application will be the proof of the Céa-type estimate for the Galerkin error (Proposition 2). Moreover, we prove that the Scott-Zhang interpolation error in a locally weighted H^1 -seminorm is locally equivalent to the Dirichlet data oscillations (Proposition 8). This will be the main tool to derive the bound $\|(1 - \mathbb{P}_\ell)g\|_{H^{1/2}(\Gamma_D)} \lesssim \text{osc}_{D, \ell}$.

3.1. Scott-Zhang projection. Analyzing the definition of \mathbb{J}_ℓ in [SZ], one sees that \mathbb{J}_ℓ can be defined locally in the following sense:

- For an element $T \in \mathcal{T}_\ell$, the value $(\mathbb{J}_\ell w)|_T$ on T depends only on the value of $w|_{\omega_{\ell, T}}$ on some element patch

$$T \subseteq \omega_{\ell, T} \subseteq \{T' \in \mathcal{T}_\ell : T' \cap T \neq \emptyset\}. \quad (31)$$

- For a boundary facet $E \in \mathcal{E}_\ell^\Gamma$, the trace of the Scott-Zhang projection $(\mathbb{J}_\ell w)|_E$ on E depends only on the trace $w|_{\omega_{\ell,E}^\Gamma}$ on some facet patch

$$E \subseteq \omega_{\ell,E}^\Gamma \subseteq \{E' \in \mathcal{E}_\ell : E' \cap E \neq \emptyset\}. \quad (32)$$

- In case of a Dirichlet facet $E \in \mathcal{E}_\ell^D$, one may choose $\omega_{\ell,E}^\Gamma \subseteq \bar{\Gamma}_D$.

Moreover, \mathbb{J}_ℓ is defined in a way that the following projection properties hold:

- $\mathbb{J}_\ell W_\ell = W_\ell$ for all $W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$,
- $(\mathbb{J}_\ell w)|_\Gamma = w|_\Gamma$ for all $w \in H^1(\Omega)$ and $W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ with $w|_\Gamma = W_\ell|_\Gamma$,
- $(\mathbb{J}_\ell w)|_{\Gamma_D} = w|_{\Gamma_D}$ for all $w \in H^1(\Omega)$ and $W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ with $w|_{\Gamma_D} = W_\ell|_{\Gamma_D}$,

i.e. the projection \mathbb{J}_ℓ preserves discrete (Dirichlet) boundary data. Finally, \mathbb{J}_ℓ satisfies the following (local) stability property

$$\|\nabla(1 - \mathbb{J}_\ell)w\|_{L^2(T)} \leq C_{sz} \|\nabla w\|_{L^2(\omega_{\ell,T})} \quad \text{for all } w \in H^1(\Omega) \quad (33)$$

and (local) first-order approximation property

$$\|(1 - \mathbb{J}_\ell)w\|_{L^2(T)} \leq C_{sz} \|h_\ell \nabla w\|_{L^2(\omega_{\ell,T})} \quad \text{for all } w \in H^1(\Omega) \quad (34)$$

where $C_{sz} > 0$ depends only on shape regularity of \mathcal{T}_ℓ , cf. [SZ]. Here, $h_\ell \in L^\infty(\Omega)$ denotes the local mesh-width function defined by $h_\ell|_T = |T|^{1/d}$ for all $T \in \mathcal{T}_\ell$. Moreover, since the overlap of the patches is controlled in terms of shape regularity, the integration domains in (33)–(34) can be replaced by Ω , i.e. (33)–(34) hold also globally.

3.2. Scott-Zhang projection onto discrete trace spaces. We stress that \mathbb{J}_ℓ induces operators

$$\mathbb{J}_\ell^\Gamma : L^2(\Gamma) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell^\Gamma) \quad \text{and} \quad \mathbb{J}_\ell^D : L^2(\Gamma_D) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell^D) \quad (35)$$

in the sense of $\mathbb{J}_\ell^\Gamma(w|_\Gamma) = (\mathbb{J}_\ell w)|_\Gamma$ and $\mathbb{J}_\ell^D(w|_{\Gamma_D}) = (\mathbb{J}_\ell^\Gamma(w|_\Gamma))|_{\Gamma_D}$ for all $w \in H^1(\Omega)$. We will thus not distinguish these operators notationally. Arguing as in [SZ], for $\gamma \in \{\Gamma, \Gamma_D, \Gamma_N\}$, one sees that \mathbb{J}_ℓ satisfies even (local) L^2 -stability

$$\|(1 - \mathbb{J}_\ell)w\|_{L^2(E)} \leq C_{sz} \|w\|_{L^2(\omega_{\ell,E}^\Gamma)} \quad \text{for all } w \in L^2(\gamma), \quad (36)$$

(local) H^1 -stability

$$\|(1 - \mathbb{J}_\ell)w\|_{H^1(E)} \leq C_{sz} \|\nabla_\Gamma w\|_{L^2(\omega_{\ell,E}^\Gamma)} \quad \text{for all } w \in H^1(\gamma), \quad (37)$$

as well as a (local) first-order approximation property

$$\|(1 - \mathbb{J}_\ell)w\|_{L^2(E)} \leq C_{sz} \|h_\ell \nabla_\Gamma w\|_{L^2(\omega_{\ell,E}^\Gamma)} \quad \text{for all } w \in H^1(\gamma). \quad (38)$$

Here, $\nabla_\Gamma(\cdot)$ denotes again the surface gradient, and $h_\ell \in L^\infty(\Gamma_D)$ denotes the local mesh-width function restricted to Γ_D . According to shape regularity of \mathcal{T}_ℓ , the integration domains in (36)–(38) can be replaced by γ , i.e. (36)–(38) hold also globally on γ .

By standard interpolation arguments applied to (36)–(37), one obtains stability

$$\|(1 - \mathbb{J}_\ell)w\|_{H^{1/2}(\gamma)} \leq C_{sz} \|w\|_{H^{1/2}(\gamma)} \quad \text{for all } w \in H^{1/2}(\gamma) \quad (39)$$

in the trace norm. Moreover, it is proved in [KOP, Theorem 3] that the Scott-Zhang projection satisfies

$$\|(1 - \mathbb{J}_\ell)w\|_{H^{1/2}(\gamma)} \leq C_{sz} \min_{W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell|_\gamma)} \|h_\ell^{1/2} \nabla_\Gamma(w - W_\ell)\|_{L^2(\gamma)} \quad \text{for all } w \in H^{1/2}(\gamma). \quad (40)$$

Throughout, the constant $C_{sz} > 0$ then depends only on shape regularity of \mathcal{T}_ℓ and on $\gamma \in \{\Gamma, \Gamma_D, \Gamma_N\}$.

3.3. Proof of Céa lemma (Proposition 2). According to weak formulation (2) and Galerkin formulation (5), we have the Galerkin orthogonality relation

$$\langle \nabla(u - U_\ell), \nabla V_\ell \rangle_\Omega = 0 \quad \text{for all } V_\ell \in \mathcal{S}_D^p(\mathcal{T}_\ell).$$

Let $\mathcal{L} : H^{1/2}(\Gamma) \rightarrow H^1(\Omega)$ be a lifting operator. Let $\widehat{g}, \widehat{g}_\ell \in H^{1/2}(\Gamma)$ denote arbitrary extensions of $g = u|_{\Gamma_D}$ resp. $\mathbb{P}_\ell g = U_\ell|_{\Gamma_D}$. Note that $(\mathbb{J}_\ell \mathcal{L} \mathbb{J}_\ell \widehat{g})|_{\Gamma_D} = (\mathbb{J}_\ell u)|_{\Gamma_D}$ as well as $(\mathbb{J}_\ell \mathcal{L} \mathbb{J}_\ell \widehat{g}_\ell)|_{\Gamma_D} = U_\ell|_{\Gamma_D}$. For arbitrary $V_\ell \in \mathcal{S}_D^p(\mathcal{T}_\ell)$, we thus have $U_\ell - (V_\ell + \mathbb{J}_\ell \mathcal{L} \mathbb{J}_\ell \widehat{g}_\ell) \in \mathcal{S}_D^p(\mathcal{T}_\ell)$, whence

$$\|\nabla(u - U_\ell)\|_{L^2(\Omega)}^2 = \langle \nabla(u - U_\ell), \nabla(u - (V_\ell + \mathbb{J}_\ell \mathcal{L} \mathbb{J}_\ell \widehat{g}_\ell)) \rangle_\Omega$$

according to the Galerkin orthogonality. Therefore, the Cauchy inequality proves

$$\|\nabla(u - U_\ell)\|_{L^2(\Omega)} \leq \min_{V_\ell \in \mathcal{S}_D^p(\mathcal{T}_\ell)} \|\nabla(u - (V_\ell + \mathbb{J}_\ell \mathcal{L} \mathbb{J}_\ell \widehat{g}_\ell))\|_{L^2(\Omega)}.$$

We now plug-in $V_\ell = \mathbb{J}_\ell u - \mathbb{J}_\ell \mathcal{L} \mathbb{J}_\ell \widehat{g} \in \mathcal{S}_D^p(\mathcal{T}_\ell)$ and use stability of \mathbb{J}_ℓ and \mathcal{L} to see

$$\begin{aligned} \|\nabla(u - U_\ell)\|_{L^2(\Omega)} &\leq \|\nabla(u - \mathbb{J}_\ell u + \mathbb{J}_\ell \mathcal{L} \mathbb{J}_\ell (\widehat{g} - \widehat{g}_\ell))\|_{L^2(\Omega)} \\ &\lesssim \|\nabla(u - \mathbb{J}_\ell u)\|_{L^2(\Omega)} + \|\widehat{g} - \widehat{g}_\ell\|_{H^{1/2}(\Gamma)}. \end{aligned}$$

Since the extensions $\widehat{g}, \widehat{g}_\ell$ of g and $\mathbb{P}_\ell g$ were arbitrary, we obtain

$$\|\nabla(u - U_\ell)\|_{L^2(\Omega)} \lesssim \|\nabla(u - \mathbb{J}_\ell u)\|_{L^2(\Omega)} + \|(1 - \mathbb{P}_\ell)g\|_{H^{1/2}(\Gamma_D)}.$$

According to the projection property $\mathbb{J}_\ell W_\ell = W_\ell$ for $W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ and H^1 -stability (33), it holds that

$$\|\nabla(u - \mathbb{J}_\ell u)\|_{L^2(\Omega)} = \min_{W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)} \|\nabla(1 - \mathbb{J}_\ell)(u - W_\ell)\|_{L^2(\Omega)} \lesssim \min_{W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)} \|\nabla(u - W_\ell)\|_{L^2(\Omega)}.$$

The same argument for \mathbb{P}_ℓ with stability on $H^{1/2}(\Gamma)$ gives

$$\|(1 - \mathbb{P}_\ell)g\|_{H^{1/2}(\Gamma_D)} = \min_{W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)} \|(1 - \mathbb{P}_\ell)(g - W_\ell|_{\Gamma_D})\|_{H^{1/2}(\Gamma_D)} \lesssim \min_{W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)} \|g - W_\ell|_{\Gamma_D}\|_{H^{1/2}(\Gamma_D)}.$$

Combining the last three estimates, we infer

$$\|\nabla(u - U_\ell)\|_{L^2(\Omega)} + \|(1 - \mathbb{P}_\ell)g\|_{H^{1/2}(\Gamma_D)} \lesssim \min_{W_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)} (\|\nabla(u - W_\ell)\|_{L^2(\Omega)} + \|g - W_\ell|_{\Gamma_D}\|_{H^{1/2}(\Gamma_D)}).$$

Finally, the Rellich compactness theorem implies norm equivalence $\|\cdot\|_{H^1(\Omega)} \simeq \|\nabla(\cdot)\|_{L^2(\Omega)} + \|(\cdot)|_{\Gamma_D}\|_{H^{1/2}(\Gamma_D)}$ on $H^1(\Omega)$. This concludes the proof. \square

3.4. Scott-Zhang projection and Dirichlet data oscillations. We stress that the newest vertex bisection algorithm guarantees that only finitely many shapes of elements $T \in \{T \in \mathcal{T}_\star : \mathcal{T}_\star \in \mathbb{T}\}$ can occur. In particular, only finitely many shapes of patches occur. Further details are found in [V, Chapter 4] as well as in [S08, T]. This observation will be used in the proof of the following lemma.

Proposition 8. Let $\Pi_\ell : L^2(\Gamma_D) \rightarrow \mathcal{P}^{p-1}(\mathcal{E}_\ell^D)$ denote the $L^2(\Gamma_D)$ -projection. Then,

$$\|(1 - \Pi_\ell)\nabla_\Gamma g\|_{L^2(E)} \leq \|\nabla_\Gamma(1 - \mathbb{J}_\ell)g\|_{L^2(E)} \leq C_{\text{dir}}\|(1 - \Pi_\ell)\nabla_\Gamma g\|_{L^2(\omega_{\ell,E}^\Gamma)} \text{ for all } E \in \mathcal{E}_\ell^D \quad (41)$$

and, in particular,

$$\text{osc}_{D,\ell} \leq \|h_\ell^{1/2}\nabla_\Gamma(1 - \mathbb{J}_\ell)g\|_{L^2(\Gamma_D)} \leq C_{\text{dir}} \text{osc}_{D,\ell} \quad (42)$$

The constant $C_{\text{dir}} \geq 1$ depends only on Γ_D , the polynomial degree p , the initial triangulation \mathcal{T}_0 , and the use of newest vertex bisection to obtain $\mathcal{T}_\ell \in \mathbb{T}$, but not on g .

Proof. Since Π_ℓ is the piecewise L^2 -projection, the lower bound in (41)–(42) is obvious. To verify the upper bound, we argue by contradiction and assume that the upper bound in (41) is wrong for each constant $C > 0$. For $n \in \mathbb{N}$, we thus find some $\tilde{g}_n \in H^1(\Gamma)$ such that

$$\|\nabla_\Gamma(1 - \mathbb{J}_\ell)\tilde{g}_n\|_{L^2(E)} > n \|(1 - \Pi_\ell)\nabla_\Gamma\tilde{g}_n\|_{L^2(\omega_{\ell,E}^\Gamma)}. \quad (43)$$

Let $\mathbb{Q}_{\ell,E} : H^1(\omega_{\ell,E}^\Gamma) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})$ denote the H^1 -orthogonal projection on the patch $\omega_{\ell,E}^\Gamma$ and define $\bar{g}_n = (1 - \mathbb{Q}_{\ell,E})\tilde{g}_n$. Since the value of $\mathbb{J}_\ell v$ on E depends only on the values of v on $\omega_{\ell,E}^\Gamma$, the projection property of \mathbb{J}_ℓ reveals $(1 - \mathbb{J}_\ell)\mathbb{Q}_{\ell,E}\tilde{g}_n = 0$ on E . Moreover, $\nabla_\Gamma\mathbb{Q}_{\ell,E}\tilde{g}_n \in \mathcal{P}^{p-1}(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})$ so that $(1 - \Pi_\ell)\nabla_\Gamma\mathbb{Q}_{\ell,E}\tilde{g}_n = 0$ on $\omega_{\ell,E}^\Gamma$. From the orthogonal decomposition $\tilde{g}_n = \mathbb{Q}_{\ell,E}\tilde{g}_n + \bar{g}_n$, we thus see $\|\nabla_\Gamma(1 - \mathbb{J}_\ell)\bar{g}_n\|_{L^2(E)} = \|\nabla_\Gamma(1 - \mathbb{J}_\ell)\tilde{g}_n\|_{L^2(E)}$ and $\|(1 - \Pi_\ell)\nabla_\Gamma\bar{g}_n\|_{L^2(\omega_{\ell,E}^\Gamma)} = \|(1 - \Pi_\ell)\nabla_\Gamma\tilde{g}_n\|_{L^2(\omega_{\ell,E}^\Gamma)}$. In particular, we observe $\bar{g}_n \neq 0$ from (43) so that we may define $g_n := \bar{g}_n / \|\bar{g}_n\|_{H^1(\omega_{\ell,E}^\Gamma)}$. This definition guarantees

$$\|g_n\|_{H^1(\omega_{\ell,E}^\Gamma)} = 1 \quad \text{and} \quad g_n \in \mathcal{S}^p(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})^\perp, \quad (44)$$

where orthogonality is understood with respect to the $H^1(\omega_{\ell,E}^\Gamma)$ -scalar product. Moreover, it holds that

$$\|(1 - \Pi_\ell)\nabla_\Gamma g_n\|_{L^2(\omega_{\ell,E}^\Gamma)} < \frac{1}{n} \|\nabla_\Gamma(1 - \mathbb{J}_\ell)g_n\|_{L^2(E)} \lesssim \frac{1}{n} \|\nabla_\Gamma g_n\|_{L^2(\omega_{\ell,E}^\Gamma)} \xrightarrow{n \rightarrow \infty} 0 \quad (45)$$

due to the construction of g_n and local H^1 -stability of $\mathbb{J}_\ell : H^1(\Gamma_D) \rightarrow H^1(\Gamma_D)$.

First, (45) implies that $\|\Pi_\ell \nabla_\Gamma g_n\|_{L^2(\omega_{\ell,E}^\Gamma)} \leq C < \infty$ is uniformly bounded as $n \rightarrow \infty$. Since $\Pi_\ell \nabla_\Gamma g_n \in \mathcal{P}^{p-1}(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})$ belongs to a finite dimensional space, we may apply the Bolzano-Weierstrass theorem to extract a convergent subsequence. Without loss of generality, we may thus assume

$$\Pi_\ell \nabla_\Gamma g_n \xrightarrow{n \rightarrow \infty} \Phi_\ell \in \mathcal{P}^{p-1}(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma}) \quad \text{in strong } L^2\text{-sense.} \quad (46)$$

Second, this and (45) prove L^2 -convergence of $\nabla_\Gamma g_n$ to Φ_ℓ ,

$$\|\nabla_\Gamma g_n - \Phi_\ell\|_{L^2(\omega_{\ell,E}^\Gamma)} \leq \|(1 - \Pi_\ell)\nabla_\Gamma g_n\|_{L^2(\omega_{\ell,E}^\Gamma)} + \|\Pi_\ell \nabla_\Gamma g_n - \Phi_\ell\|_{L^2(\omega_{\ell,E}^\Gamma)} \xrightarrow{n \rightarrow \infty} 0. \quad (47)$$

Third, orthogonality (44) implies $\int_{\omega_{\ell,E}^\Gamma} g_n d\Gamma = 0$ if we consider the constant function $1 \in \mathcal{S}^p(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})$. Therefore, the Friedrichs inequality and (47) predict uniform boundedness $\|g_n\|_{H^1(\omega_{\ell,E}^\Gamma)} \lesssim \|\nabla_\Gamma g_n\|_{L^2(\omega_{\ell,E}^\Gamma)} \leq C < \infty$ as $n \rightarrow \infty$. According to weak compactness in Hilbert spaces, we may thus extract a weakly convergent subsequence. Without loss of generality, we may thus assume

$$g_n \xrightarrow{n \rightarrow \infty} g_\infty \in H^1(\omega_{\ell,E}^\Gamma) \quad \text{in weak } H^1\text{-sense.} \quad (48)$$

Fourth, the combination of (47) and (48) implies $\nabla_\Gamma g_\infty = \Phi_\ell$. This follows from the fact that $\|\Phi_\ell - \nabla_\Gamma(\cdot)\|_{L^2(\omega_{\ell,E}^\Gamma)}$ is convex and continuous, whence weakly lower semicontinuous on $H^1(\omega_{\ell,E}^\Gamma)$, i.e. $\|\Phi_\ell - \nabla_\Gamma g_\infty\|_{L^2(\omega_{\ell,E}^\Gamma)} \leq \liminf_n \|\Phi_\ell - \nabla_\Gamma g_n\|_{L^2(\omega_{\ell,E}^\Gamma)} = 0$.

Fifth, the Rellich compactness theorem proves that the convergence in (48) does also hold in strong L^2 -sense. Together with (47) and $\Phi_\ell = \nabla_\Gamma g_\infty$, we now observe strong H^1 -convergence

$$\|g_\infty - g_n\|_{H^1(\omega_{\ell,E}^\Gamma)}^2 = \|g_\infty - g_n\|_{L^2(\omega_{\ell,E}^\Gamma)}^2 + \|\Phi_\ell - \nabla_\Gamma g_n\|_{L^2(\omega_{\ell,E}^\Gamma)}^2 \xrightarrow{n \rightarrow \infty} 0,$$

whence $\|g_\infty\|_{H^1(\omega_{\ell,E}^\Gamma)} = 1$ as well as $g_\infty \in \mathcal{S}^p(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})^\perp$ according to (44).

On the other hand, $\nabla_\Gamma g_\infty = \Phi_\ell \in \mathcal{P}^{p-1}(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})$ implies $g_\infty \in \mathcal{S}^p(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})$. This yields $g_\infty \in \mathcal{S}^p(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma}) \cap \mathcal{S}^p(\mathcal{E}_\ell|_{\omega_{\ell,E}^\Gamma})^\perp = \{0\}$ and contradicts $\|g_\infty\|_{H^1(\omega_{\ell,E}^\Gamma)}^2 = 1$.

This contradiction proves the upper bound in (41). A standard scaling argument verifies that the constant $C_{\text{dir}} > 0$ does only depend on the shape of $\omega_{\ell,E}^\Gamma$ but not on the diameter. As stated above, newest vertex bisection guarantees that only finitely many shapes of patches $\omega_{\ell,E}^\Gamma$ may occur, i.e. $C_{\text{dir}} > 0$ depends only on \mathcal{T}_0 and the use of newest vertex bisection. Summing (41) over all Dirichlet facets, we see

$$\begin{aligned} \text{osc}_{D,\ell}^2 &= \|h_\ell^{1/2}(1 - \Pi_\ell)\nabla_\Gamma g\|_{L^2(\Gamma_D)}^2 \leq \|h_\ell^{1/2}\nabla_\Gamma(1 - \mathbb{J}_\ell)g\|_{L^2(\Gamma_D)}^2 \\ &\lesssim \sum_{E \in \mathcal{E}_\ell^D} h_\ell|_E \|(1 - \Pi_\ell)\nabla_\Gamma g\|_{L^2(\omega_{\ell,E}^\Gamma)}^2 \\ &\lesssim \|h_\ell^{1/2}(1 - \Pi_\ell)\nabla_\Gamma g\|_{L^2(\Gamma_D)}^2, \end{aligned}$$

where the final estimate holds due to uniform shape regularity. \square

Corollary 9. *It holds $\|(1 - \mathbb{P}_\ell)g\|_{H^{1/2}(\Gamma_D)} \leq C_{\text{osc}} \text{osc}_{D,\ell}$, where $C_{\text{osc}} > 0$ depends on Γ_D , the polynomial degree $p \geq 1$, stability $C_{\text{stab}} > 0$, the initial mesh \mathcal{T}_0 , and the use of newest vertex bisection.*

Proof. By use of the projection property and stability of \mathbb{P}_ℓ , one sees $\|(1 - \mathbb{P}_\ell)g\|_{H^{1/2}(\Gamma_D)} = \|(1 - \mathbb{P}_\ell)(1 - \mathbb{J}_\ell)g\|_{H^{1/2}(\Gamma_D)} \lesssim \|(1 - \mathbb{J}_\ell)g\|_{H^{1/2}(\Gamma_D)}$. The approximation estimate (40) and Proposition 8 conclude $\|(1 - \mathbb{J}_\ell)g\|_{H^{1/2}(\Gamma_D)} \lesssim \|h_\ell^{1/2}\nabla_\Gamma(1 - \mathbb{J}_\ell)g\|_{L^2(\Gamma_D)} \simeq \text{osc}_{D,\ell}$. \square

3.5. Proof of reliability and efficiency (Proposition 3). We consider a continuous auxiliary problem

$$\begin{aligned} -\Delta w &= 0 && \text{in } \Omega, \\ w &= (1 - \mathbb{P}_\ell)g && \text{on } \Gamma_D, \\ \partial_n w &= 0 && \text{on } \Gamma_N, \end{aligned} \tag{49}$$

with unique solution $w \in H^1(\Omega)$. We then have norm equivalence $\|w\|_{H^1(\Omega)} \simeq \|(1 - \mathbb{P}_\ell)g\|_{H^{1/2}(\Gamma_D)}$ as well as $u - U_\ell - w \in H_D^1(\Omega)$. From this, we obtain

$$\|u - U_\ell\|_{H^1(\Omega)} \lesssim \|\nabla(u - U_\ell - w)\|_{L^2(\Omega)} + \|(1 - \mathbb{P}_\ell)g\|_{H^{1/2}(\Gamma_D)}.$$

The first term on the right-hand side can be handled as for homogeneous Dirichlet data, i.e. use of the Galerkin orthogonality combined with approximation estimates for a Clément-type quasi-interpolation operator (e.g. the Scott-Zhang projection). This leads to

$$\|\nabla(u - U_\ell - w)\|_{L^2(\Omega)} \lesssim \varrho_\ell$$

Details are found e.g. in [BCD]. The $H^{1/2}(\Gamma_D)$ -norm is dominated by the Dirichlet data oscillations $\text{osc}_{D,\ell}$, see Corollary 9.

By use of bubble functions and local scaling arguments, one obtains the estimates

$$|T|^{2/d} \|f + \Delta U_\ell\|_{L^2(T)}^2 \lesssim \|\nabla(u - U_\ell)\|_{L^2(T)}^2 + \text{osc}_{\mathcal{T},\ell}(T)^2 + \text{osc}_{N,\ell}(\partial T \cap \Gamma_N),$$

$$|T|^{1/d} \|[\partial_n U_\ell]\|_{L^2(E \cap \Omega)}^2 \lesssim \|\nabla(u - U_\ell)\|_{L^2(\Omega_{\ell,E})}^2 + \text{osc}_{\mathcal{T},\ell}(\omega_{\ell,E})^2,$$

$$|T|^{1/d} \|\phi - \partial_n U_\ell\|_{L^2(E \cap \Gamma_N)}^2 \lesssim \|\nabla(u - U_\ell)\|_{L^2(\Omega_{\ell,E})}^2 + \text{osc}_{\mathcal{T},\ell}(\omega_{\ell,E})^2 + \text{osc}_{N,\ell}(E \cap \Gamma_N)^2,$$

where $\Omega_{\ell,E} = T^+ \cup T^-$ denotes the facet patch of $T_+ \cap T_- = E \in \mathcal{E}_\ell$. Details are found e.g. in [AO, V]. Summing these estimates over all elements, one obtains the efficiency estimate (16). \square

4. CONVERGENCE

In this section, we aim to prove Theorem 4. Our proof of the convergence theorem relies on the estimator reduction principle from [AFP], i.e. we verify that the error estimator is contractive up to some zero sequence.

4.1. Estimator reduction estimate. Note that the estimator $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ can be localized over elements via

$$\eta_\ell^2 = \sum_{T \in \mathcal{T}_\ell} \eta_\ell(T)^2 \quad \text{with} \quad \eta_\ell(T)^2 = \varrho_\ell(T)^2 + |T|^{1/d} \|(1 - \Pi_\ell) \nabla_\Gamma g\|_{L^2(\partial T \cap \Gamma_D)}^2 \quad (50)$$

with $\Pi_\ell : L^2(\Gamma_D) \rightarrow \mathcal{P}^{p-1}(\mathcal{E}_\ell^D)$ the (even \mathcal{E}_ℓ^D -piecewise) $L^2(\Gamma_D)$ -orthogonal projection.

Lemma 10 (modified marking implies Dörfler marking). *For $0 < \theta_1, \theta_2, \vartheta < 1$ in Algorithm 1, there is some parameter $0 < \theta < 1$ such that the error estimator $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ satisfies*

$$\theta \eta_\ell^2 \leq \sum_{T \in \mathcal{M}_\ell} \eta_\ell(T)^2, \quad (51)$$

and all elements $T \in \mathcal{M}_\ell$ are refined by at least one bisection.

Proof. First, assume $\text{osc}_{D,\ell}^2 \leq \vartheta \varrho_\ell^2$ and let $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell$ satisfy (11). Then,

$$\theta_1(\varrho_\ell^2 + \text{osc}_{D,\ell}^2) \leq \theta_1(1 + \vartheta)\varrho_\ell^2 \leq (1 + \vartheta) \sum_{T \in \mathcal{M}_\ell} \varrho_\ell(T)^2.$$

Therefore, the Dörfler marking (51) holds with $\theta \leq \theta_1(1 + \vartheta)^{-1}$.

Second, assume $\text{osc}_{D,\ell}^2 > \vartheta \varrho_\ell^2$ and let $\mathcal{M}_\ell^D \subseteq \mathcal{E}_\ell^D$ satisfy (12). Then,

$$\theta_2(\varrho_\ell^2 + \text{osc}_{D,\ell}^2) \leq \theta_2(1 + \vartheta^{-1})\text{osc}_{D,\ell}^2 \leq (1 + \vartheta^{-1}) \sum_{E \in \mathcal{M}_\ell^D} \text{osc}_{D,\ell}(E)^2.$$

Therefore, the Dörfler marking (51) holds with $\theta \leq \theta_2(1 + \vartheta^{-1})^{-1}$, and all elements which have some facet $E \in \mathcal{M}_\ell^D$ are refined. \square

Proposition 11 (estimator reduction). *Let $\mathcal{T}_\star = \mathbf{refine}(\mathcal{T}_\ell)$ be an arbitrary refinement of \mathcal{T}_ℓ and $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell \setminus \mathcal{T}_\star$ a subset of the refined elements which satisfies the Dörfler marking (51) for some $0 < \theta < 1$. Then,*

$$\eta_\star^2 \leq q_{\text{red}} \eta_\ell^2 + C_{\text{red}} \|\nabla(U_{\ell+1} - U_\ell)\|_{L^2(\Omega)}^2 \quad (52)$$

with certain constants $0 < q_{\text{red}} < 1$ and $C_{\text{red}} > 0$ which depend only on the parameter $0 < \theta < 1$, shape regularity of \mathcal{T}_\star , and the polynomial degree $p \geq 1$.

Sketch of proof. For the sake of completeness, we include the idea of the proof of (52) although our proof is only a minor extension of the proof from [CKNS, Cor. 3.4, Proof of Theorem 4.1], where all details are found. First, we employ a triangle inequality and the Young inequality to see for arbitrary $\delta > 0$

$$\begin{aligned} \eta_\star^2 \leq (1 + \delta) & \left(\sum_{T' \in \mathcal{T}_\star} |T'|^{2/d} \|f + \Delta U_\ell\|_{L^2(T')}^2 + |T'|^{1/d} \|[\partial_n U_\ell]\|_{L^2(\partial T' \cap \Omega)}^2 \right. \\ & \quad \left. + |T'|^{1/d} \|\phi - \partial_n U_\ell\|_{L^2(\partial T' \cap \Gamma_N)}^2 + |T'|^{1/d} \|(1 - \Pi_\star) \nabla_\Gamma g\|_{L^2(\partial T' \cap \Gamma_D)}^2 \right) \\ & + (1 + \delta^{-1}) \left(\sum_{T' \in \mathcal{T}_\star} |T'|^{2/d} \|\Delta(U_\star - U_\ell)\|_{L^2(T')}^2 + |T'|^{1/d} \|[\partial_n(U_\star - U_\ell)]\|_{L^2(\partial T' \cap \Omega)}^2 \right. \\ & \quad \left. + |T'|^{1/d} \|\partial_n(U_\star - U_\ell)\|_{L^2(\partial T' \cap \Gamma_N)}^2 \right). \end{aligned}$$

A scaling argument proves that the second bracket is bounded by $C \|\nabla(U_\star - U_\ell)\|_{L^2(\Omega)}^2$, where $C > 0$ depends only on shape regularity of \mathcal{T}_\star . The first contribution of the first bracket is estimated as follows

$$\begin{aligned} \sum_{T' \in \mathcal{T}_\star} |T'|^{2/d} \|f + \Delta U_\ell\|_{L^2(T')}^2 &= \sum_{T' \in \mathcal{T}_\star \cap \mathcal{T}_\ell} |T'|^{2/d} \|f + \Delta U_\ell\|_{L^2(T')}^2 \\ &\quad + \sum_{T' \in \mathcal{T}_\star \setminus \mathcal{T}_\ell} |T'|^{2/d} \|f + \Delta U_\ell\|_{L^2(T')}^2 \end{aligned}$$

Since a refined element $T \in \mathcal{T}_\ell \setminus \mathcal{T}_\star$ is the (essentially disjoint) union of its sons $T' \in \mathcal{T}_\star \setminus \mathcal{T}_\ell$ and $|T'| \leq |T|/2$, the second sum is estimated by

$$\sum_{T' \in \mathcal{T}_\star \setminus \mathcal{T}_\ell} |T'|^{2/d} \|f + \Delta U_\ell\|_{L^2(T')}^2 \leq 2^{-2/d} \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{T}_\star} |T|^{2/d} \|f + \Delta U_\ell\|_{L^2(T)}^2.$$

This yields

$$\begin{aligned} \sum_{T' \in \mathcal{T}_\star} |T'|^{2/d} \|f + \Delta U_\ell\|_{L^2(T')}^2 &\leq \sum_{T \in \mathcal{T}_\ell} |T|^{2/d} \|f + \Delta U_\ell\|_{L^2(T)}^2 \\ &\quad - (1 - 2^{-2/d}) \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{T}_\star} |T|^{2/d} \|f + \Delta U_\ell\|_{L^2(T)}^2, \end{aligned}$$

where the sums on the right-hand side only involve contributions of $\eta_\ell(T)^2$. We employ the same type of argument for the other contributions. Together with the estimate

$$\|(1 - \Pi_\star) \nabla_\Gamma g\|_{L^2(\partial T' \cap \Gamma_D)} \leq \|(1 - \Pi_\ell) \nabla_\Gamma g\|_{L^2(\partial T' \cap \Gamma_D)} \quad \text{for all } T' \in \mathcal{T}_\star$$

which follows from the fact that the L^2 -projection onto $\mathcal{P}^{p-1}(\mathcal{E}_\star^D)$ is even piecewise orthogonal, we are led to

$$\begin{aligned}
& \left(\sum_{T' \in \mathcal{T}_\star} |T'|^{2/d} \|f + \Delta U_\ell\|_{L^2(T')}^2 + |T'|^{1/d} \|[\partial_n U_\ell]\|_{L^2(\partial T' \cap \Omega)}^2 \right. \\
& \quad \left. + |T'|^{1/d} \|\phi - \partial_n U_\ell\|_{L^2(\partial T' \cap \Gamma_N)}^2 + |T'|^{1/d} \|(1 - \Pi_\star) \nabla_\Gamma g\|_{L^2(\partial T' \cap \Gamma_D)}^2 \right) \\
& \leq \sum_{T \in \mathcal{T}_\ell} \eta_\ell(T)^2 - (1 - 2^{-1/d}) \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{T}_\star} \eta_\ell(T)^2 \\
& \leq (1 - (1 - 2^{-1/d})\theta) \eta_\ell^2.
\end{aligned}$$

The final estimate follows from $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell \setminus \mathcal{T}_\star$ and the Dörfler marking (51), i.e. we subtract less. With $0 < q := 1 - (1 - 2^{-1/d})\theta$, we have thus proved

$$\eta_\star^2 \leq (1 + \delta) q \eta_\ell^2 + (1 + \delta^{-1}) C \|\nabla(U_\star - U_\ell)\|_{L^2(\Omega)}^2.$$

Finally, we choose $\delta > 0$ sufficiently small such that $0 < q_{\text{red}} := (1 + \delta) q < 1$ and define $C_{\text{red}} = (1 + \delta^{-1}) C$. \square

4.2. A priori convergence of Scott-Zhang projection. We assume that $(\mathbb{J}_{\ell+1} v)|_T = (\mathbb{J}_\ell v)|_T$ for all $T \in \mathcal{T}_\ell \cap \mathcal{T}_{\ell+1}$ with $\omega_{\ell,T} \subseteq \bigcup(\mathcal{T}_\ell \cap \mathcal{T}_{\ell+1})$ which can always be achieved by an appropriate choice of the dual basis functions in the definition of $\mathbb{J}_{\ell+1}$. In this section, we prove that under the foregoing assumptions and for arbitrary refinement, i.e. $\mathcal{T}_\ell = \text{refine}(\mathcal{T}_{\ell-1})$ for all $\ell \in \mathbb{N}$, the limit of the Scott-Zhang interpolants $\mathbb{J}_\ell v$ exists in $H^1(\Omega)$ as $\ell \rightarrow \infty$. In particular, this provides the essential ingredient to prove that, under the same assumptions, the limit of Galerkin solutions U_ℓ exists in $H^1(\Omega)$. For 2D and first-order elements $p = 1$, this result has first been proved in [FPP]. Although the proof transfers directly to the present setting, we include it for the sake of completeness.

Proposition 12. *Let $v \in H^1(\Omega)$. Then, the limit $\mathbb{J}_\infty v := \lim_\ell \mathbb{J}_\ell v$ exists in $H^1(\Omega)$ and defines a continuous linear operator $\mathbb{J}_\infty : H^1(\Omega) \rightarrow H^1(\Omega)$.*

Proof. If the limit $\mathbb{J}_\infty v$ exists for all $v \in H^1(\Omega)$, it is a consequence of the Banach-Steinhaus theorem that \mathbb{J}_∞ defines, in fact, a linear and continuous operator. Hence, it remains to prove the existence of $\mathbb{J}_\infty v$ in $H^1(\Omega)$ for fixed $v \in H^1(\Omega)$. To that end, we follow the ideas from [MSV] and define the following subsets of Ω :

$$\begin{aligned}
\Omega_\ell^0 &:= \bigcup \{T \in \mathcal{T}_\ell : \omega_\ell(T) \subseteq \bigcup(\bigcup_{i=0}^\infty \bigcap_{j=i}^\infty \mathcal{T}_j)\}, \\
\Omega_\ell &:= \bigcup \{T \in \mathcal{T}_\ell : \text{There exists } k \geq 0 \text{ s.t. } \omega_\ell(T) \text{ is at least uniformly refined in } \mathcal{T}_{\ell+k}\}, \\
\Omega_\ell^* &:= \Omega \setminus (\Omega_\ell \cup \Omega_\ell^0),
\end{aligned}$$

where $\omega_\ell(\omega) := \bigcup \{T \in \mathcal{T}_\ell : T \cap \omega \neq \emptyset\}$ denotes the patch of $\omega \subseteq \Omega$ with respect to \mathcal{T}_ℓ . In other words, Ω_ℓ^0 is the set of all elements whose patch is not refined anymore and thus stays the same in \mathcal{T}_k for all $k \geq \ell$, whereas Ω_ℓ denotes the set of elements whose patch is uniformly refined at least once after k steps. According to [MSV, Corollary 4.1] and uniform shape regularity, it holds that

$$\lim_{\ell \rightarrow \infty} \|\chi_{\Omega_\ell} h_\ell\|_{L^\infty(\Omega)} = 0 = \lim_{\ell \rightarrow \infty} \|\chi_{\omega_\ell(\Omega_\ell)} h_\ell\|_{L^\infty(\Omega)}, \quad (53)$$

where $\chi_{\Omega_\ell} : \Omega \rightarrow \{0, 1\}$ denotes the characteristic function with respect to Ω_ℓ and $\chi_{\omega_\ell(\Omega_\ell)}$ the characteristic function of the patch of Ω_ℓ . Now, let $\varepsilon > 0$ be arbitrary. Since the space $H^2(\Omega)$ is dense in $H^1(\Omega)$, we find $v_\varepsilon \in H^2(\Omega)$ such that $\|v - v_\varepsilon\|_{H^1(\Omega)} \leq \varepsilon$. Due to local approximation and stability properties of \mathbb{J}_ℓ (see (33) and (34)), we obtain

$$\|(1 - \mathbb{J}_\ell)v\|_{H^1(\Omega_\ell)} \lesssim \|(1 - \mathbb{J}_\ell)v_\varepsilon\|_{H^1(\Omega_\ell)} + \varepsilon \leq \|h_\ell D^2 v_\varepsilon\|_{L^2(\omega_\ell(\Omega_\ell))} + \varepsilon,$$

where the last estimate is a consequence of the Bramble-Hilbert lemma. By use of (53), we may choose $\ell_0 \in \mathbb{N}$ sufficiently large to guarantee the estimate $\|h_\ell D^2 v_\varepsilon\|_{L^2(\omega_\ell(\Omega_\ell))} \leq \|h_\ell\|_{L^\infty(\omega_\ell(\Omega_\ell))} \|D^2 v_\varepsilon\|_{L^2(\Omega)} \leq \varepsilon$ for all $\ell \geq \ell_0$. Then, there holds

$$\|(1 - \mathbb{J}_\ell)v\|_{H^1(\Omega_\ell)} \lesssim \varepsilon \quad \text{for all } \ell \geq \ell_0. \quad (54)$$

According to [MSV, Proposition 4.2], it holds $\lim_\ell |\Omega_\ell^*| = 0$. This provides the existence of $\ell_1 \in \mathbb{N}$ such that

$$\|v\|_{H^1(\omega_\ell(\Omega_\ell^*))} \leq \varepsilon \quad \text{for all } \ell \geq \ell_1 \quad (55)$$

due to the non-concentration of Lebesgue functions and uniform shape regularity, i.e. $|\omega_\ell(\Omega_\ell^*)| \lesssim |\Omega_\ell^*|$. With these preparations, we finally aim at proving that $\mathbb{J}_\ell v$ is a Cauchy sequence in $H^1(\Omega)$: Let $\ell \geq \max\{\ell_0, \ell_1\}$ and $k \geq 0$ be arbitrary. First, we use that for any $T \in \mathcal{T}_\ell$, $(\mathbb{J}_\ell v)|_T$ depends only on $v|_{\omega_\ell(T)}$ because $\omega_{\ell,T} \subseteq \omega_\ell(T)$. Then, by definition of Ω_ℓ^0 and our assumption on the definition of \mathbb{J}_ℓ and $\mathbb{J}_{\ell+k}$ on $\mathcal{T}_\ell \cap \mathcal{T}_{\ell+k}$, we obtain

$$\|\mathbb{J}_\ell v - \mathbb{J}_{\ell+k} v\|_{H^1(\Omega_\ell^0)} = 0. \quad (56)$$

Second, due to the local stability of \mathbb{J}_ℓ and (55), there holds

$$\begin{aligned} \|\mathbb{J}_\ell v - \mathbb{J}_{\ell+k} v\|_{H^1(\Omega_\ell^*)} &\leq \|\mathbb{J}_\ell v\|_{H^1(\Omega_\ell^*)} + \|\mathbb{J}_{\ell+k} v\|_{H^1(\Omega_\ell^*)} \\ &\lesssim \|v\|_{H^1(\omega_\ell(\Omega_\ell^*))} + \|v\|_{H^1(\omega_{\ell+k}(\Omega_\ell^*))} \\ &\leq 2\|v\|_{H^1(\omega_\ell(\Omega_\ell^*))} \leq 2\varepsilon. \end{aligned} \quad (57)$$

Third, we proceed by exploiting (54). We have

$$\|\mathbb{J}_\ell v - \mathbb{J}_{\ell+k} v\|_{H^1(\Omega_\ell)} \leq \|\mathbb{J}_\ell v - v\|_{H^1(\Omega_\ell)} + \|v - \mathbb{J}_{\ell+k} v\|_{H^1(\Omega_\ell)} \lesssim \varepsilon. \quad (58)$$

Combining the estimates from (56)–(58), we conclude $\|\mathbb{J}_\ell v - \mathbb{J}_{\ell+k} v\|_{H^1(\Omega)} \lesssim \varepsilon$, i.e. $(\mathbb{J}_\ell v)$ is a Cauchy sequence in $H^1(\Omega)$ and hence convergent. \square

Corollary 13. *Under the assumptions of Proposition 12, the limit $g_\infty := \lim_\ell \mathbb{J}_\ell g$ exists in $H^{1/2}(\Gamma_D)$.*

Proof. Let $\widehat{g} \in H^{1/2}(\Gamma)$ denote an arbitrary extension of g . With some lifting operator \mathcal{L} , we define $v := \mathcal{L}\widehat{g}$ and note that $(\mathbb{J}_\ell v)|_{\Gamma_D} = (\mathbb{J}_\ell \widehat{g})|_{\Gamma_D} = \mathbb{J}_\ell g$. Since $\mathbb{J}_\infty v = \lim_\ell \mathbb{J}_\ell v$ exists in $H^1(\Omega)$, we obtain

$$\|(\mathbb{J}_\infty v)|_{\Gamma_D} - \mathbb{J}_\ell g\|_{H^{1/2}(\Gamma_D)} \leq \|(\mathbb{J}_\infty v)|_\Gamma - \mathbb{J}_\ell \widehat{g}\|_{H^{1/2}(\Gamma)} \leq \|\mathbb{J}_\infty v - \mathbb{J}_\ell v\|_{H^1(\Omega)} \xrightarrow{\ell \rightarrow \infty} 0.$$

This concludes the proof with $(\mathbb{J}_\infty v)|_{\Gamma_D} =: g_\infty$. \square

4.3. A priori convergence of orthogonal projections. In this subsection, we recall an early observation from [BV, Lemma 6.1] which will be applied several times. We stress that the original proof of [BV] is based on the orthogonal projection. However, the argument also works for (possibly nonlinear) projections with $P_\ell P_k = P_\ell$ for $\ell \leq k$ which satisfy a

Céa-type quasi-optimality. Since the Scott-Zhang projection satisfies $\mathbb{J}_\ell \mathbb{J}_k \neq \mathbb{J}_\ell$, in general, Proposition 12 is *not* a consequence of such an abstract result.

Lemma 14. *Let H be a Hilbert space and X_ℓ be a sequence of closed subspaces with $X_\ell \subseteq X_{\ell+1}$ for all $\ell \geq 0$. Let $P_\ell : H \rightarrow X_\ell$ denote the H -orthogonal projection onto X_ℓ . Then, for each $x \in H$, the limit $x_\infty := \lim_{\ell \rightarrow \infty} P_\ell x$ exists in H . \square*

Since the discrete trace spaces $\mathcal{S}^p(\mathcal{E}_\ell^D)$ are finite dimensional and hence closed subspaces of $H^{1/2}(\Gamma_D)$, the lemma immediately applies to the $H^{1/2}(\Gamma_D)$ -orthogonal projection.

Corollary 15. *Let $P_\ell : H^{1/2}(\Gamma_D) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell^D)$ denote the $H^{1/2}(\Gamma_D)$ -orthogonal projection. Then, the limit $g_\infty := \lim_{\ell \rightarrow \infty} P_\ell g$ exists in $H^{1/2}(\Gamma_D)$. \square*

Corollary 16. *Let $\pi_\ell : L^2(\Gamma_D) \rightarrow \mathcal{S}^1(\mathcal{E}_\ell^D)$ denote the $L^2(\Gamma_D)$ -orthogonal projection. Then, the limit $g_\infty := \lim_{\ell \rightarrow \infty} \pi_\ell g$ exists weakly in $H^1(\Gamma_D)$ and strongly in $H^\alpha(\Gamma_D)$ for all $0 \leq \alpha < 1$.*

Proof. According to Lemma 14, the limit $g_\infty = \lim_{\ell} \pi_\ell g$ exists strongly in $L^2(\Gamma_D)$. Moreover and according to [KPP, Theorem 8], the π_ℓ are uniformly stable in $H^1(\Gamma_D)$, since we use newest vertex bisection. Hence, the sequence $(\pi_\ell g)$ is uniformly bounded in $H^1(\Gamma_D)$ and thus admits a weakly convergent subsequence $(\pi_{\ell_k} g)$ with weak limit $\tilde{g}_\infty \in H^1(\Gamma_D)$, where weak convergence is understood in $H^1(\Gamma_D)$. Since the inclusion $H^1(\Gamma_D) \subset L^2(\Gamma_D)$ is compact, the sequence $(\pi_{\ell_k} g)$ converges strongly to \tilde{g}_∞ in $L^2(\Gamma_D)$. From uniqueness of limits, we conclude $\tilde{g}_\infty = g_\infty$. Iterating this argument, we see that each subsequence of $(\pi_\ell g)$ contains a subsequence which converges weakly to g_∞ in $H^1(\Gamma_D)$. This proves that the entire sequence converges weakly to g_∞ in $H^1(\Gamma_D)$. Strong convergence in $H^\alpha(\Gamma_D)$ follows by compact inclusion $H^1(\Gamma_D) \subset H^\alpha(\Gamma_D)$ for all $0 \leq \alpha < 1$. \square

4.4. A priori convergence of Galerkin solutions. We now show that the limit of Galerkin solutions U_ℓ exists as $\ell \rightarrow \infty$ provided that the meshes are nested, i.e. $\mathcal{T}_{\ell+1} = \text{refine}(\mathcal{T}_\ell)$.

Proposition 17. *Under Assumption (17) that $g_\infty := \lim_{\ell} \mathbb{P}_\ell g$ exists in $H^{1/2}(\Gamma)$, also the limit $U_\infty := \lim_{\ell} U_\ell$ of Galerkin solutions exists in $H^1(\Omega)$.*

Proof. We consider the continuous auxiliary problem

$$\begin{aligned} -\Delta w_\ell &= 0 && \text{in } \Omega, \\ w_\ell &= \mathbb{P}_\ell g && \text{on } \Gamma_D, \\ \partial_n w_\ell &= 0 && \text{on } \Gamma_N. \end{aligned}$$

Let $w_\ell \in H^1(\Omega)$ be the unique (weak) solution and note that the trace $\widehat{g}_\ell := w_\ell|_\Gamma \in H^{1/2}(\Gamma)$ provides an extension of $\mathbb{P}_\ell g$ with

$$\|\widehat{g}_\ell\|_{H^{1/2}(\Gamma)} \leq \|w_\ell\|_{H^1(\Omega)} \lesssim \|\mathbb{P}_\ell g\|_{H^{1/2}(\Gamma_D)} \leq \|\widehat{g}_\ell\|_{H^{1/2}(\Gamma)}.$$

For arbitrary $k, \ell \in \mathbb{N}$, the same type of arguments proves

$$\|\widehat{g}_\ell - \widehat{g}_k\|_{H^{1/2}(\Gamma)} \simeq \|(\mathbb{P}_\ell - \mathbb{P}_k)g\|_{H^{1/2}(\Gamma_D)}.$$

According to Assumption (17), $(\mathbb{P}_\ell g)$ is a Cauchy sequence in $H^{1/2}(\Gamma_D)$. Therefore, (\widehat{g}_ℓ) is a Cauchy sequence in $H^{1/2}(\Gamma)$, whence convergent with limit $\widehat{g}_\infty \in H^{1/2}(\Gamma)$. Next, note that

$(\mathbb{J}_\ell \mathcal{L} \hat{g}_\ell)|_{\Gamma_D} = \mathbb{P}_\ell g$, where $\mathcal{L} : H^{1/2}(\Gamma) \rightarrow H^1(\Omega)$ denotes some lifting operator. Therefore, $\tilde{U}_\ell := U_\ell - \mathbb{J}_\ell \mathcal{L} \hat{g}_\ell \in \mathcal{S}_D^p(\mathcal{T}_\ell)$ is the unique solution of the variational formulation

$$\langle \nabla \tilde{U}_\ell, \nabla V_\ell \rangle_\Omega = \langle \nabla u, \nabla V_\ell \rangle_\Omega - \langle \nabla \mathbb{J}_\ell \mathcal{L} \hat{g}_\ell, \nabla V_\ell \rangle_\Omega \quad \text{for all } V_\ell \in \mathcal{S}_D^p(\mathcal{T}_\ell). \quad (59)$$

Finally, we need to show that \tilde{U}_ℓ and $\mathbb{J}_\ell \mathcal{L} \hat{g}_\ell$ are convergent to conclude convergence of $U_\ell = \tilde{U}_\ell + \mathbb{J}_\ell \mathcal{L} \hat{g}_\ell$.

With convergence of (\hat{g}_ℓ) to \hat{g}_∞ and Proposition 12, we obtain

$$\begin{aligned} \|\mathbb{J}_\ell \mathcal{L} \hat{g}_\ell - \mathbb{J}_\infty \mathcal{L} \hat{g}_\infty\|_{H^1(\Omega)} &\leq \|\mathbb{J}_\ell(\mathcal{L} \hat{g}_\ell - \mathcal{L} \hat{g}_\infty)\|_{H^1(\Omega)} + \|\mathbb{J}_\ell \mathcal{L} \hat{g}_\infty - \mathbb{J}_\infty \mathcal{L} \hat{g}_\infty\|_{H^1(\Omega)} \\ &\lesssim \|\hat{g}_\ell - \hat{g}_\infty\|_{H^{1/2}(\Gamma)} + \|\mathbb{J}_\ell \mathcal{L} \hat{g}_\infty - \mathbb{J}_\infty \mathcal{L} \hat{g}_\infty\|_{H^1(\Omega)} \xrightarrow{\ell \rightarrow \infty} 0. \end{aligned}$$

This proves convergence of $\mathbb{J}_\ell \mathcal{L} \hat{g}_\ell$ to $\mathbb{J}_\infty \mathcal{L} \hat{g}_\infty$ as $\ell \rightarrow \infty$. To see convergence of \tilde{U}_ℓ , let $\tilde{U}_{\ell,\infty} \in \mathcal{S}_D^p(\mathcal{T}_\ell)$ be the unique solution of the discrete auxiliary problem

$$\langle \nabla \tilde{U}_{\ell,\infty}, \nabla V_\ell \rangle_\Omega = \langle \nabla u, \nabla V_\ell \rangle_\Omega - \langle \nabla \mathbb{J}_\infty \mathcal{L} \hat{g}_\infty, \nabla V_\ell \rangle_\Omega \quad \text{for all } V_\ell \in \mathcal{S}_D^p(\mathcal{T}_\ell). \quad (60)$$

Due to the nestedness of the ansatz spaces $\mathcal{S}_D^p(\mathcal{T}_\ell)$, Lemma 14 predicts a priori convergence $\tilde{U}_{\ell,\infty} \xrightarrow{\ell \rightarrow \infty} \tilde{U}_\infty \in H^1(\Omega_D)$. With the stability of (59) and (60), we obtain

$$\|\nabla(\tilde{U}_{\ell,\infty} - \tilde{U}_\ell)\|_{L^2(\Omega)} \lesssim \|\mathbb{J}_\ell \mathcal{L} \hat{g}_\ell - \mathbb{J}_\infty \mathcal{L} \hat{g}_\infty\|_{H^1(\Omega)} \xrightarrow{\ell \rightarrow \infty} 0,$$

and therefore $\tilde{U}_\ell \xrightarrow{\ell \rightarrow \infty} \tilde{U}_\infty$ in $H^1(\Omega_D)$.

Finally, we now conclude

$$U_\ell = \tilde{U}_\ell + \mathbb{J}_\ell \mathcal{L} \hat{g}_\ell \xrightarrow{\ell \rightarrow \infty} \tilde{U}_\infty + \mathbb{J}_\infty \mathcal{L} \hat{g}_\infty =: U_\infty \in H^1(\Omega),$$

which concludes the proof. \square

4.5. Proof of convergence theorem (Theorem 4). (i) Since the limit $U_\infty = \lim_\ell U_\ell$ exists in $H^1(\Omega)$, we infer $\lim_\ell \|\nabla(U_{\ell+1} - U_\ell)\|_{L^2(\Omega)} = 0$. In view of this and Lemma 10, the estimator reduction estimate (52) takes the form

$$\eta_{\ell+1}^2 \leq q_{\text{red}} \eta_\ell^2 + \alpha_\ell \quad \text{for all } \ell \geq 0$$

with some non-negative $\alpha_\ell \geq 0$ such that $\lim_\ell \alpha_\ell = 0$, i.e. the estimator is contractive up to a non-negative zero sequence. It is a consequence of elementary calculus that $\lim_\ell \eta_\ell = 0$, see e.g. [AFP, Lemma 2.3]. Finally, reliability $\|u - U_\ell\|_{H^1(\Omega)} \lesssim \eta_\ell$ thus concludes the proof.

(ii) The verification of Assumption (17) is done in Corollary 13 for the Scott-Zhang projection, Corollary 15 for the $H^{1/2}(\Gamma_D)$ -orthogonal projection, and Corollary 16 for the $L^2(\Gamma_D)$ -orthogonal projection. \square

5. CONTRACTION

In principle, the convergence rate of $\lim_\ell U_\ell = u$ from Theorem 4 could be slow. Moreover, Theorem 4 restricts the Dirichlet projection \mathbb{P}_ℓ by Assumption (17). In this section, we aim to show linear convergence for some quasi-error quantity $\Delta_\ell \simeq \eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ with respect to the step ℓ of Algorithm 1 and independently of the projection \mathbb{P}_ℓ chosen. The essential observation is that the marking step in Algorithm 1 is in some sense independent of the \mathbb{P}_ℓ chosen.

5.1. Implicit Dörfler marking. Let $\tilde{U}_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ be a Galerkin solution of (5) with different Dirichlet data $\tilde{U}_\ell = P_\ell g$ on Γ_D , where $P_\ell : H^{1/2}(\Gamma_D) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell^D)$ is a uniformly stable projection onto $\mathcal{S}^p(\mathcal{E}_\ell^D)$ in the sense of (13). Let $\tilde{\eta}_\ell^2 = \tilde{\varrho}_\ell^2 + \text{osc}_{D,\ell}^2$ be the associated error estimator. In the following, we prove that marking in Algorithm 1 with $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ and sufficiently small $0 < \vartheta < 1$ implicitly implies the simple Dörfler marking (51) for $\tilde{\eta}_\ell$.

Lemma 18 (local equivalence of error estimators for different projections). *For arbitrary $\mathcal{U}_\ell \subseteq \mathcal{T}_\ell$, it holds that*

$$C_{\text{eq}}^{-1} \sum_{T \in \mathcal{U}_\ell} \varrho_\ell(T)^2 \leq \sum_{T \in \mathcal{U}_\ell} \tilde{\varrho}_\ell(T)^2 + \text{osc}_{D,\ell}^2 \quad \text{and} \quad C_{\text{eq}}^{-1} \sum_{T \in \mathcal{U}_\ell} \tilde{\varrho}_\ell(T)^2 \leq \sum_{T \in \mathcal{U}_\ell} \varrho_\ell(T)^2 + \text{osc}_{D,\ell}^2. \quad (61)$$

The constant $C_{\text{eq}} > 1$ depends only on shape regularity of \mathcal{T}_ℓ and on $C_{\text{stab}} > 0$. In particular, this implies equivalence

$$(C_{\text{eq}} + 1)^{-1} \eta_\ell^2 \leq \tilde{\eta}_\ell^2 \leq (C_{\text{eq}} + 1) \eta_\ell^2. \quad (62)$$

Proof. Arguing as for the estimator reduction, it follows from the triangle inequality and scaling arguments that

$$\varrho_\ell(T)^2 \lesssim \tilde{\varrho}_\ell(T)^2 + \|\nabla(U_\ell - \tilde{U}_\ell)\|_{L^2(\omega_\ell(T))}^2 \quad \text{for all } T \in \mathcal{T}_\ell,$$

where $\omega_\ell(T) = \bigcup \{T' \in \mathcal{T}_\ell : T' \cap T \neq \emptyset\}$ denotes the element patch of T . Consequently, a rough estimate gives

$$\sum_{T \in \mathcal{U}_\ell} \varrho_\ell(T)^2 \lesssim \sum_{T \in \mathcal{U}_\ell} \tilde{\varrho}_\ell(T)^2 + \|\nabla(U_\ell - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 \quad \text{for all } \mathcal{U}_\ell \subseteq \mathcal{T}_\ell.$$

Recall the Galerkin orthogonality

$$\langle \nabla(U_\ell - \tilde{U}_\ell), \nabla V_\ell \rangle = \langle \nabla(u - \tilde{U}_\ell), \nabla V_\ell \rangle - \langle \nabla(u - U_\ell), \nabla V_\ell \rangle = 0 \quad \text{for all } V_\ell \in \mathcal{S}_D^p(\mathcal{T}_\ell).$$

Let $\hat{g} \in H^{1/2}(\Gamma)$ be an arbitrary extension of $(U_\ell - \tilde{U}_\ell)|_{\Gamma_D} = (\mathbb{P}_\ell - P_\ell)g \in H^{1/2}(\Gamma_D)$. We choose the test function $V_\ell = (U_\ell - \tilde{U}_\ell) - \mathbb{J}_\ell \mathcal{L} \hat{g} \in \mathcal{S}_D^1(\mathcal{T}_\ell)$ to see

$$\|\nabla(U_\ell - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 = \langle \nabla(U_\ell - \tilde{U}_\ell), \nabla \mathbb{J}_\ell \mathcal{L} \hat{g} \rangle_\Omega.$$

Stability of Scott-Zhang projection \mathbb{J}_ℓ and lifting operator \mathcal{L} thus give

$$\|\nabla(U_\ell - \tilde{U}_\ell)\|_{L^2(\Omega)} \leq \|\nabla \mathbb{J}_\ell \mathcal{L} \hat{g}\|_{L^2(\Omega)} \lesssim \|\hat{g}\|_{H^{1/2}(\Gamma)}.$$

Since \hat{g} was an arbitrary extension of $(\mathbb{P}_\ell - P_\ell)g$, we end up with

$$\begin{aligned} \|\nabla(U_\ell - \tilde{U}_\ell)\|_{L^2(\Omega)} &\lesssim \|(\mathbb{P}_\ell - P_\ell)g\|_{H^{1/2}(\Gamma_D)} \leq \|(\mathbb{P}_\ell - 1)g\|_{H^{1/2}(\Gamma_D)} + \|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)} \\ &\lesssim \text{osc}_{D,\ell}, \end{aligned}$$

where we have used Corollary 9. This proves the first estimate in (61), and the second follows with the same arguments. \square

The following lemma is the main reason, why we stick with Stevenson's modified Dörfler marking (11)–(12) instead of simple Dörfler marking (51).

Lemma 19 (modified Dörfler marking implies Dörfler marking for different projection). *For arbitrary $0 < \theta_1, \theta_2 < 1$ and sufficiently small $0 < \vartheta < 1$, there is some $0 < \theta < 1$ such that the marking criterion (11)–(12) for $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ implies the Dörfler marking*

$$\theta \tilde{\eta}_\ell^2 \leq \sum_{T \in \mathcal{M}_\ell} \tilde{\eta}_\ell(T)^2 \quad (63)$$

for $\tilde{\eta}_\ell^2 = \tilde{\varrho}_\ell^2 + \text{osc}_{D,\ell}^2$. The parameter $0 < \theta < 1$ depends on $0 < \theta_1, \theta_2, \vartheta < 1$ and on $C_{\text{eq}} > 0$.

Proof. We argue as in the proof of Lemma 10. First, assume $\text{osc}_{D,\ell}^2 \leq \vartheta \varrho_\ell^2$ and let $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell$ satisfy (11). According to Lemma 18, we see

$$\begin{aligned} \theta_1 \eta_\ell^2 &\leq \theta_1 (1 + \vartheta) \varrho_\ell^2 \leq (1 + \vartheta) \sum_{T \in \mathcal{M}_\ell} \varrho_\ell(T)^2 \leq C_{\text{eq}} (1 + \vartheta) \left(\sum_{T \in \mathcal{M}_\ell} \tilde{\varrho}_\ell(T)^2 + \text{osc}_{D,\ell}^2 \right) \\ &\leq C_{\text{eq}} (1 + \vartheta) \left(\sum_{T \in \mathcal{M}_\ell} \tilde{\varrho}_\ell(T)^2 + \vartheta \varrho_\ell^2 \right). \end{aligned}$$

This proves

$$(\theta_1 C_{\text{eq}}^{-1} (1 + \vartheta)^{-1} - \vartheta) \eta_\ell^2 \leq \sum_{T \in \mathcal{M}_\ell} \tilde{\varrho}_\ell(T)^2.$$

Together with $(C_{\text{eq}} + 1)^{-1} \tilde{\eta}_\ell^2 \leq \eta_\ell^2$, we thus obtain the Dörfler marking (63) with $0 < \theta \leq (C_{\text{eq}} + 1)^{-1} (\theta_1 C_{\text{eq}}^{-1} (1 + \vartheta)^{-1} - \vartheta) < 1$, provided that $0 < \vartheta < 1$ is sufficiently small compared to $0 < \theta_1 < 1$.

Second, assume $\text{osc}_{D,\ell}^2 > \vartheta \varrho_\ell^2$ and let $\mathcal{M}_\ell^D \subseteq \mathcal{E}_\ell^D$ satisfy (12). Then,

$$\theta_2 \eta_\ell^2 \leq \theta_2 (1 + \vartheta^{-1}) \text{osc}_{D,\ell}^2 \leq (1 + \vartheta^{-1}) \sum_{E \in \mathcal{M}_\ell^D} \text{osc}_{D,\ell}(E)^2 \leq (1 + \vartheta^{-1}) \sum_{T \in \mathcal{M}_\ell} \tilde{\eta}_\ell(T)^2,$$

where $\mathcal{M}_\ell = \{T \in \mathcal{T}_\ell : \exists E \in \mathcal{M}_\ell^D \text{ } E \subset \partial T\}$ is defined in step (iv) of Algorithm 1. As before $(C_{\text{eq}} + 1)^{-1} \tilde{\eta}_\ell^2 \leq \eta_\ell^2$ thus proves (51) with $0 < \theta \leq C_{\text{eq}}^{-1} \theta_2 (1 + \vartheta^{-1})^{-1} < 1$. \square

5.2. Quasi-Pythagoras theorem. To prove Theorem 5, we consider a *theoretical* auxiliary problem: Throughout the remainder of Section 5, $\tilde{U}_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ denotes the Galerkin solution of (5) with Dirichlet data $\tilde{U}_\ell = P_\ell g$ on Γ_D , where $P_\ell : H^{1/2}(\Gamma_D) \rightarrow \mathcal{S}^p(\mathcal{E}_\ell^D)$ denotes the $H^{1/2}(\Gamma_D)$ -orthogonal projection. Associated with \tilde{U}_ℓ is the error estimator $\tilde{\eta}_\ell^2 = \tilde{\varrho}_\ell^2 + \text{osc}_{D,\ell}^2$, where $\tilde{\varrho}_\ell$ is defined in (10) with U_ℓ replaced by \tilde{U}_ℓ .

Recall that the aforementioned statements of Section 3 and Section 4 hold for *any* uniformly $H^{1/2}(\Gamma_D)$ -stable projection \mathbb{P}_ℓ and thus apply to $\tilde{\eta}_\ell^2 = \tilde{\varrho}_\ell^2 + \text{osc}_{D,\ell}^2$. We shall need reliability $\|u - \tilde{U}_\ell\|_{H^1(\Omega)}^2 \lesssim \tilde{\eta}_\ell^2$ as well as the estimator reduction (52) from Proposition 11 for $\tilde{\eta}_\ell^2$, which is a consequence of Lemma 19. Our concept of proof of Theorem 5 goes back to [CKNS, Proof of Theorem 4.1]. Therein, however, the proof relies on the Pythagoras theorem $\|\nabla(u - U_\ell)\|_{L^2(\Omega)}^2 = \|\nabla(u - U_{\ell+1})\|_{L^2(\Omega)}^2 + \|\nabla(U_{\ell+1} - U_\ell)\|_{L^2(\Omega)}^2$ which does *not* hold in case of inhomogeneous Dirichlet data and $\mathbb{P}_\ell g \neq \mathbb{P}_{\ell+1} g$, in general. Instead, we rely on a perturbed Pythagoras theorem which will be used for the auxiliary problem.

Lemma 20 (quasi-Pythagoras theorem). *Let $\mathcal{T}_\star = \mathbf{refine}(\mathcal{T}_\ell)$ be an arbitrary refinement of \mathcal{T}_ℓ with the associated auxiliary solution $\tilde{U}_\star \in \mathcal{S}^p(\mathcal{T}_\star)$, where $\tilde{U}_\star = P_\star g$ on Γ_D . Then,*

$$(1 - \alpha) \|\nabla(u - \tilde{U}_\star)\|_{L^2(\Omega)}^2 \leq \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 - \|\nabla(\tilde{U}_\star - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + \alpha^{-1} C_{\text{pyth}} \|(P_\star - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 \quad (64)$$

for all $\alpha > 0$. The constant $C_{\text{pyth}} > 0$ depends only on the shape regularity of $\sigma(\mathcal{T}_\ell)$ and $\sigma(\mathcal{T}_\star)$ and on Ω and Γ_D .

Proof. We recall the Galerkin orthogonality

$$\langle \nabla(u - \tilde{U}_\star), \nabla V_\star \rangle_\Omega = 0 \quad \text{for all } V_\star \in \mathcal{S}_D^p(\mathcal{T}_\star).$$

Now, let $\tilde{U}_\star^\ell \in \mathcal{S}^p(\mathcal{T}_\star)$ be the unique Galerkin solution of (5) with $\tilde{U}_\star^\ell|_{\Gamma_D} = P_\ell g$. We use the Galerkin orthogonality with $V_\star = \tilde{U}_\star^\ell - \tilde{U}_\ell \in \mathcal{S}_D^p(\mathcal{T}_\star)$. This and the Young inequality allow to estimate the L^2 -scalar product

$$\begin{aligned} 2|\langle \nabla(u - \tilde{U}_\star), \nabla(\tilde{U}_\star - \tilde{U}_\ell) \rangle_\Omega| &= 2|\langle \nabla(u - \tilde{U}_\star), \nabla(\tilde{U}_\star - \tilde{U}_\star^\ell) \rangle_\Omega| \\ &\leq \alpha \|\nabla(u - \tilde{U}_\star)\|_{L^2(\Omega)}^2 + \alpha^{-1} \|\nabla(\tilde{U}_\star - \tilde{U}_\star^\ell)\|_{L^2(\Omega)}^2 \end{aligned}$$

for all $\alpha > 0$. To estimate the second contribution on the right-hand side, we choose an arbitrary extension $\hat{g} \in H^{1/2}(\Gamma)$ of $(P_\star - P_\ell)g \in H^{1/2}(\Gamma_D)$. Then, we use the test function $V_\star = (\tilde{U}_\star - \tilde{U}_\star^\ell) - \mathbb{J}_\star \mathcal{L} \hat{g} \in \mathcal{S}_D^p(\mathcal{T}_\star)$, where $\mathcal{L} : H^{1/2}(\Gamma) \rightarrow H^1(\Omega)$ again denotes a lifting operator. Recall that the choice of the Scott-Zhang projection \mathbb{J}_\star guarantees that this function has zero Dirichlet data on Γ_D since $\hat{g}|_{\Gamma_D} = (P_\star - P_\ell)g \in \mathcal{S}^p(\mathcal{E}_\star^D)$. Now, the Galerkin orthogonalities for $\tilde{U}_\star, \tilde{U}_\star^\ell \in \mathcal{S}^p(\mathcal{T}_\star)$ yield

$$0 = \langle \nabla(u - \tilde{U}_\star^\ell), \nabla V_\star \rangle_\Omega - \langle \nabla(u - \tilde{U}_\star), \nabla V_\star \rangle_\Omega = \langle \nabla(\tilde{U}_\star - \tilde{U}_\star^\ell), \nabla V_\star \rangle_\Omega.$$

By the above choice of $V_\star \in \mathcal{S}_D^p(\mathcal{T}_\star)$ and stability of \mathbb{J}_\star and \mathcal{L} , this yields

$$\|\nabla(\tilde{U}_\star - \tilde{U}_\star^\ell)\|_{L^2(\Omega)}^2 = \langle \nabla(\tilde{U}_\star - \tilde{U}_\star^\ell), \nabla \mathbb{J}_\star \mathcal{L} \hat{g} \rangle_\Omega \lesssim \|\nabla(\tilde{U}_\star - \tilde{U}_\star^\ell)\|_{L^2(\Omega)} \|\hat{g}\|_{H^{1/2}(\Gamma)}.$$

Since \hat{g} was an arbitrary extension of $(P_\star - P_\ell)g \in H^{1/2}(\Gamma_D)$ to $H^{1/2}(\Gamma)$, this yields

$$\|\nabla(\tilde{U}_\star - \tilde{U}_\star^\ell)\|_{L^2(\Omega)} \lesssim \|(P_\star - P_\ell)g\|_{H^{1/2}(\Gamma_D)}.$$

So far, we have thus derived

$$2|\langle \nabla(u - \tilde{U}_\star), \nabla(\tilde{U}_\star - \tilde{U}_\ell) \rangle_\Omega| \leq \alpha \|\nabla(u - \tilde{U}_\star)\|_{L^2(\Omega)}^2 + \alpha^{-1} C_{\text{pyth}} \|(P_\star - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2,$$

To verify (64), we use the identity

$$\begin{aligned} \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 &= \|\nabla((u - \tilde{U}_\star) + (\tilde{U}_\star - \tilde{U}_\ell))\|_{L^2(\Omega)}^2 \\ &= \|\nabla(u - \tilde{U}_\star)\|_{L^2(\Omega)}^2 + 2\langle \nabla(u - \tilde{U}_\star), \nabla(\tilde{U}_\star - \tilde{U}_\ell) \rangle_\Omega + \|\nabla(\tilde{U}_\star - \tilde{U}_\ell)\|_{L^2(\Omega)}^2. \end{aligned}$$

Rearranging the terms accordingly and use of the estimate for the scalar product, we conclude the proof. \square

5.3. Proof of contraction theorem (Theorem 5). Using the quasi-Pythagoras theorem (64) with $\mathcal{T}_\star = \mathcal{T}_{\ell+1}$, we see

$$(1 - \alpha) \|\nabla(u - \tilde{U}_{\ell+1})\|_{L^2(\Omega)}^2 \leq \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 - \|\nabla(\tilde{U}_{\ell+1} - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 \\ + \alpha^{-1} C_{\text{pyth}} \|(P_{\ell+1} - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2.$$

The use of the $H^{1/2}(\Gamma_D)$ -orthogonal projection provides the orthogonality relation

$$\|(1 - P_{\ell+1})g\|_{H^{1/2}(\Gamma_D)}^2 + \|(P_{\ell+1} - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 = \|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2.$$

Combining the last two estimates, we obtain

$$(1 - \alpha) \|\nabla(u - \tilde{U}_{\ell+1})\|_{L^2(\Omega)}^2 + \alpha^{-1} C_{\text{pyth}} \|(1 - P_{\ell+1})g\|_{H^{1/2}(\Gamma_D)}^2 \\ \leq \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + \alpha^{-1} C_{\text{pyth}} \|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 - \|\nabla(\tilde{U}_{\ell+1} - \tilde{U}_\ell)\|_{L^2(\Omega)}^2.$$

Applying Lemma 19, we see that Algorithm 1 for $\eta_\ell^2 = \varrho_\ell^2 + \text{osc}_{D,\ell}^2$ implicitly implies the Dörfler marking (63) (resp. (51)) for $\tilde{\eta}_\ell^2 = \tilde{\varrho}_\ell^2 + \text{osc}_{D,\ell}^2$. Therefore, the estimator reduction (52) of Proposition 11 applies to the auxiliary problem and provides

$$\tilde{\eta}_{\ell+1}^2 \leq q_{\text{red}} \tilde{\eta}_\ell^2 + C_{\text{red}} \|\nabla(\tilde{U}_{\ell+1} - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 \quad \text{for all } \ell \geq 0.$$

Now, we add the last two estimates to see, for $\beta > 0$,

$$(1 - \alpha) \|\nabla(u - \tilde{U}_{\ell+1})\|_{L^2(\Omega)}^2 + \alpha^{-1} C_{\text{pyth}} \|(1 - P_{\ell+1})g\|_{H^{1/2}(\Gamma_D)}^2 + \beta \tilde{\eta}_{\ell+1}^2 \\ \leq \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + \alpha^{-1} C_{\text{pyth}} \|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 + \beta q_{\text{red}} \tilde{\eta}_\ell^2 \\ + (\beta C_{\text{red}} - 1) \|\nabla(\tilde{U}_{\ell+1} - \tilde{U}_\ell)\|_{L^2(\Omega)}^2.$$

We choose $\beta > 0$ sufficiently small to guarantee $\beta C_{\text{red}} - 1 \leq 0$, i.e. the last term on the right-hand side of the last estimate can be omitted. Then, we use the reliability $\|u - \tilde{U}_\ell\|_{H^1(\Omega)}^2 \lesssim \tilde{\eta}_\ell^2$ and the estimate $\|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 \lesssim \text{osc}_{D,\ell}^2 \leq \tilde{\eta}_\ell^2$ from Corollary 9 in the form

$$\|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + \|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 \leq C \tilde{\eta}_\ell^2$$

to see, for arbitrary $\gamma, \delta > 0$

$$(1 - \alpha) \|\nabla(u - \tilde{U}_{\ell+1})\|_{L^2(\Omega)}^2 + \alpha^{-1} C_{\text{pyth}} \|(1 - P_{\ell+1})g\|_{H^{1/2}(\Gamma_D)}^2 + \beta \tilde{\eta}_{\ell+1}^2 \\ \leq \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + \alpha^{-1} C_{\text{pyth}} \|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 + \beta q_{\text{red}} \tilde{\eta}_\ell^2 \\ \leq (1 - \gamma \beta C^{-1}) \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + (1 - \delta \beta C^{-1}) \alpha^{-1} C_{\text{pyth}} \|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 \\ + \beta (q_{\text{red}} + \gamma + \delta \alpha^{-1} C_{\text{pyth}}) \tilde{\eta}_\ell^2.$$

For $0 < \alpha < 1$, we may now rearrange this estimate to end up with

$$\|\nabla(u - \tilde{U}_{\ell+1})\|_{L^2(\Omega)}^2 + \frac{C_{\text{pyth}}}{\alpha(1 - \alpha)} \|(1 - P_{\ell+1})g\|_{H^{1/2}(\Gamma_D)}^2 + \frac{\beta}{1 - \alpha} \tilde{\eta}_{\ell+1}^2 \\ \leq \frac{1 - \gamma \beta C^{-1}}{1 - \alpha} \|\nabla(u - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + (1 - \delta \beta C^{-1}) \frac{C_{\text{pyth}}}{\alpha(1 - \alpha)} \|(1 - P_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 \\ + (q_{\text{red}} + \gamma + \delta \alpha^{-1} C_{\text{pyth}}) \frac{\beta}{1 - \alpha} \tilde{\eta}_\ell^2.$$

It remains to choose the free constants $0 < \alpha, \gamma, \delta < 1$, whereas $\beta > 0$ has already been fixed:

- First, choose $0 < \gamma < 1$ sufficiently small to guarantee $0 < q_{\text{red}} + \gamma < 1$ and $0 < \gamma\beta C^{-1} < 1$.
- Second, choose $0 < \alpha < 1$ sufficiently small such that $0 < (1 - \gamma\beta C^{-1})/(1 - \alpha) < 1$.
- Third, choose $\delta > 0$ sufficiently small with $q_{\text{red}} + \gamma + \delta\alpha^{-1}C_{\text{pyth}} < 1$.

With $\mu := \beta/(1 - \alpha)$, $\lambda := \alpha^{-1}C_{\text{pyth}}/(1 - \alpha)$, and $0 < \kappa < 1$ the maximal contraction constant of the three contributions, we end up with the contraction estimate (20).

It thus only remains to prove equivalence (21): According to the definition of Δ_ℓ in (20), we have equivalence $\Delta_\ell \simeq \tilde{\eta}_\ell^2$. Finally, Lemma 18 implies $\tilde{\eta}_\ell^2 \simeq \eta_\ell^2$ and concludes the proof. \square

Remark. For $\mathbb{P}_\ell = P_\ell$ the $H^{1/2}(\Gamma_D)$ -orthogonal projection, the proof reveals that Theorem 5 holds without any restriction on the adaptivity parameters and even for simple Dörfler marking (51). \square

6. QUASI-OPTIMALITY

In this section, we aim to prove Theorem 6–7. In some sense, the heart of the matter of the quasi-optimality analysis is the discrete local reliability of Proposition 21. This is, however, only proved for discrete Dirichlet data obtained by the Scott-Zhang projection. We therefore consider this as an auxiliary problem: Let $\tilde{U}_\ell \in \mathcal{S}^p(\mathcal{T}_\ell)$ denote the Galerkin solution of (5) with respect to the Scott-Zhang projection, i.e. $\tilde{U}_\ell = \mathbb{J}_\ell g$ on Γ_D . Finally and as above, $\tilde{\eta}_\ell^2 = \tilde{\varrho}_\ell^2 + \text{osc}_{D,\ell}^2$ denotes the error estimator for this auxiliary problem. Although the discrete local reliability of $\tilde{\eta}_\ell$ does not imply discrete local reliability of the error estimator η_ℓ for the primal problem, we will see that nevertheless discrete local reliability of an equivalent error estimator is sufficient for quasi-optimality.

6.1. Optimality of Dörfler marking. Throughout, we assume that the Scott-Zhang projections are chosen with respect to the assumptions of Section 4.2.

Proposition 21 (discrete local reliability for Scott-Zhang projection). *Let $\mathcal{T}_\star = \text{refine}(\mathcal{T}_\ell)$ be an arbitrary refinement of \mathcal{T}_ℓ and $\tilde{U}_\star \in \mathcal{S}^p(\mathcal{T}_\star)$ the corresponding Galerkin solution (5) with $\tilde{U}_\star = \mathbb{J}_\star g$ on Γ_D . Then, there is a set $\mathcal{R}_\ell \subseteq \mathcal{T}_\ell$ which contains the refined elements, $\mathcal{T}_\ell \setminus \mathcal{T}_\star \subseteq \mathcal{R}_\ell$ such that*

$$\|\tilde{U}_\star - \tilde{U}_\ell\|_{H^1(\Omega)} \leq C_{\text{dlr}} \sum_{T \in \mathcal{R}_\ell} \tilde{\eta}_\ell(T)^2 \quad \text{and} \quad \#\mathcal{R}_\ell \leq C_{\text{ref}} \#(\mathcal{T}_\ell \setminus \mathcal{T}_\star). \quad (65)$$

The constants $C_{\text{dlr}}, C_{\text{ref}} > 0$ depend only on \mathcal{T}_0 and the use of newest vertex bisection.

Proof. We consider a discrete auxiliary problem

$$\langle \nabla W_\star, \nabla V_\star \rangle_\Omega = 0 \quad \text{for all } V_\star \in \mathcal{S}_D^p(\mathcal{T}_\star)$$

with unique solution $W_\star \in \mathcal{S}^p(\mathcal{T}_\star)$ with $W_\star|_{\Gamma_D} = (\mathbb{J}_\star - \mathbb{J}_\ell)g$. Then, $(\tilde{U}_\star - \tilde{U}_\ell - W_\star) \in \mathcal{S}_D^p(\mathcal{T}_\star)$, and the H^1 -norm is bounded by the H^1 -seminorm. Moreover, arguing as in [CKNS,

Lemma 3.6], we see

$$\|\tilde{U}_\star - \tilde{U}_\ell - W_\star\|_{H^1(\Omega)}^2 \lesssim \|\nabla(\tilde{U}_\star - \tilde{U}_\ell - W_\star)\|_{L^2(\Omega)}^2 \lesssim \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{T}_\star} \tilde{\varrho}_\ell(T)^2 \leq \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{T}_\star} \tilde{\eta}_\ell(T)^2.$$

According to the triangle inequality, it thus only remains to bound $\|W_\star\|_{H^1(\Omega)}$ by $\sum_{T \in \mathcal{R}_\ell} \tilde{\eta}_\ell(T)^2$ with some appropriate $\mathcal{R}_\ell \supseteq \mathcal{T}_\ell \setminus \mathcal{T}_\star$. To that end, let $\mathcal{L} : H^{1/2}(\Gamma) \rightarrow H^1(\Omega)$ be a lifting operator and $\hat{g} \in H^{1/2}(\Gamma)$ an arbitrary extension of $(\mathbb{J}_\star - \mathbb{J}_\ell)g \in H^{1/2}(\Gamma_D)$. With $V_\star := W_\star - \mathbb{J}_\star \mathcal{L} \hat{g} \in \mathcal{S}_D^p(\mathcal{T}_\star)$, we obtain

$$\begin{aligned} \|W_\star\|_{L^2(\Omega)} &\leq \|V_\star\|_{L^2(\Omega)} + \|\mathbb{J}_\star \mathcal{L} \hat{g}\|_{L^2(\Omega)} \lesssim \|\nabla V_\star\|_{L^2(\Omega)} + \|\mathbb{J}_\star \mathcal{L} \hat{g}\|_{L^2(\Omega)} \\ &\lesssim \|\nabla W_\star\|_{L^2(\Omega)} + \|\mathbb{J}_\star \mathcal{L} \hat{g}\|_{H^1(\Omega)}. \end{aligned}$$

Moreover, the variational formulation for $W_\star \in \mathcal{S}^p(\mathcal{T}_\star)$ yields

$$0 = \langle \nabla W_\star, \nabla V_\star \rangle_\Omega = \|\nabla W_\star\|_{L^2(\Omega)}^2 - \langle \nabla W_\star, \nabla \mathbb{J}_\star \mathcal{L} \hat{g} \rangle_\Omega, \text{ whence } \|\nabla W_\star\|_{L^2(\Omega)} \leq \|\nabla \mathbb{J}_\star \mathcal{L} \hat{g}\|_{L^2(\Omega)}.$$

Combining the last two estimates, we obtain

$$\|W_\star\|_{H^1(\Omega)} \lesssim \|\mathbb{J}_\star \mathcal{L} \hat{g}\|_{H^1(\Omega)} \lesssim \|\hat{g}\|_{H^{1/2}(\Gamma)}.$$

Since \hat{g} was an arbitrary extension, this proves

$$\|W_\star\|_{H^1(\Omega)} \lesssim \|(\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{H^{1/2}(\Gamma_D)}.$$

To abbreviate the notation in the remainder of the proof, let $\mathcal{R}_\ell^D := \mathcal{E}_\ell^D \setminus \mathcal{E}_\ell^\star$ denote the refined Dirichlet facets. We define inductively

$$\omega_\ell^0 = \bigcup \mathcal{R}_\ell^D, \quad \omega_\ell^n = \bigcup \{E \in \mathcal{E}_\ell^D : E \cap \omega_\ell^{n-1} \neq \emptyset\} \quad \text{for } n \geq 1,$$

i.e. ω_ℓ^n denotes the region of the refined Dirichlet facets plus n layers of (non-refined) Dirichlet facets with respect to \mathcal{E}_ℓ^D . Note that ω_ℓ^1 is nothing but the usual patch of \mathcal{R}_ℓ^D . Due to the local definition of \mathbb{J}_ℓ and \mathbb{J}_\star , we observe

$$(\mathbb{J}_\star - \mathbb{J}_\ell)g = 0 \quad \text{on } \Gamma_D \setminus \omega_\ell^1. \quad (66)$$

Let $\zeta_{\ell,z} \in \mathcal{S}^1(\mathcal{E}_\ell^D)$ denote the hat function associated with some node $z \in \mathcal{K}_\ell^D$ of \mathcal{E}_ℓ^D . Clearly, the hat functions $\{\zeta_{\ell,z} : z \in \mathcal{K}_\ell^D\}$ provide a partition of unity $\sum_{z \in \mathcal{K}_\ell^D} \zeta_{\ell,z} = 1$ on Γ_D resp. $\sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \zeta_{\ell,z} = 1$ on ω_ℓ^1 . Exploiting (66), we see

$$\|(\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{H^{1/2}(\Gamma_D)} = \left\| \sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \zeta_{\ell,z} (\mathbb{J}_\star - \mathbb{J}_\ell)g \right\|_{H^{1/2}(\Gamma_D)}. \quad (67)$$

We now adapt the arguments of [CMS, FKMP] to our setting. Analogously to the proof of [CMS, Theorem 3.2] resp. [FKMP, Proposition 4.3], we obtain

$$\begin{aligned} \left\| \sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \zeta_{\ell,z} (\mathbb{J}_\star - \mathbb{J}_\ell)g \right\|_{H^{1/2}(\Gamma_D)}^2 &\lesssim \sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \|\zeta_{\ell,z} (\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 \\ &\lesssim \sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \|\zeta_{\ell,z} (\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{L^2(\Gamma_D)} \|\zeta_{\ell,z} (\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{H^1(\Gamma_D)}, \end{aligned}$$

where the final estimate is just the interpolation estimate. As above, let

$$\omega_{\ell,z}^0 := \{z\}, \quad \omega_{\ell,z}^n := \bigcup \{E \in \mathcal{E}_\ell^D : E \cap \omega_{\ell,z}^{n-1} \neq \emptyset\} \quad \text{for } n \geq 1,$$

i.e. $\omega_{\ell,z}^1 = \bigcup \{E \in \mathcal{E}_\ell^D : z \in E\}$ denotes the node patch of $z \in \mathcal{K}_\ell^D$ which is just the support of the hat function $\zeta_{\ell,z}$ on Γ_D . To proceed, we apply the Friedrichs inequality to the summands on the right-hand side of the estimate above and derive

$$\begin{aligned} \left\| \sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \zeta_{\ell,z}(\mathbb{J}_\star - \mathbb{J}_\ell)g \right\|_{H^{1/2}(\Gamma_D)}^2 &\lesssim \sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \text{diam}(\omega_{\ell,z}^1) \|\nabla_\Gamma(\zeta_{\ell,z}(\mathbb{J}_\star - \mathbb{J}_\ell)g)\|_{L^2(\omega_{\ell,z}^1)}^2 \\ &\simeq \sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \|h_\ell^{1/2} \nabla_\Gamma(\zeta_{\ell,z}(\mathbb{J}_\star - \mathbb{J}_\ell)g)\|_{L^2(\omega_{\ell,z}^1)}^2. \end{aligned} \quad (68)$$

Here, $h_\ell \in L^\infty(\Gamma_D)$ denotes the local mesh-width function $h_\ell|_E = |T|^{1/d}$ for $E \in \mathcal{E}_\ell^D$ and $T \in \mathcal{T}_\ell$ the unique element with $E \subset \partial T$. Formally, the constants in the Friedrichs inequality depend on the shape of $\omega_{\ell,z}^1$. Note, however, that there are only finitely many shapes of patches due to the use of newest vertex bisection. Next, we use the estimate $|\nabla_\Gamma \zeta_{\ell,z}|_E| \simeq \text{diam}(E)^{-1} \simeq h_\ell^{-1}|_E$ for $E \in \mathcal{E}_\ell^D$. This and the product rule yield

$$\begin{aligned} &\|h_\ell^{1/2} \nabla_\Gamma(\zeta_{\ell,z}(\mathbb{J}_\star - \mathbb{J}_\ell)g)\|_{L^2(\omega_{\ell,z}^1)}^2 \\ &\leq \|h_\ell^{1/2}(\nabla_\Gamma \zeta_{\ell,z})(\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^1)}^2 + \|h_\ell^{1/2} \nabla_\Gamma(\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^1)}^2 \\ &\lesssim \|h_\ell^{-1/2}(\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^1)}^2 + \|h_\ell^{1/2} \nabla_\Gamma(\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^1)}^2 \\ &= \|h_\ell^{-1/2} \mathbb{J}_\star(1 - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^1)}^2 + \|h_\ell^{1/2} \nabla_\Gamma \mathbb{J}_\star(1 - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^1)}^2. \end{aligned}$$

Finally, the local stability of \mathbb{J}_\star and the local approximation property of \mathbb{J}_ℓ yield

$$\begin{aligned} \|h_\ell^{1/2} \nabla_\Gamma(\zeta_{\ell,z}(\mathbb{J}_\star - \mathbb{J}_\ell)g)\|_{L^2(\omega_{\ell,z}^1)}^2 &\lesssim \|h_\ell^{-1/2}(1 - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^2)}^2 + \|h_\ell^{1/2} \nabla_\Gamma(1 - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^2)}^2 \\ &\lesssim \|h_\ell^{1/2} \nabla_\Gamma(1 - \mathbb{J}_\ell)g\|_{L^2(\omega_{\ell,z}^3)}^2 \\ &\lesssim \|h_\ell^{1/2}(1 - \Pi_\ell) \nabla_\Gamma g\|_{L^2(\omega_{\ell,z}^4)}^2, \end{aligned} \quad (69)$$

where we have finally used Estimate (41) of Proposition 8. Now, let $\tilde{\mathcal{R}}_\ell^D := \{E \in \mathcal{E}_\ell^D : E \subseteq \omega_\ell^5\}$ denote the set of Dirichlet facets which lie in ω_ℓ^5 and note that $\#\tilde{\mathcal{R}}_\ell^D \simeq \#\mathcal{R}_\ell^D \lesssim \#(\mathcal{T}_\ell \setminus \mathcal{T}_\star)$ up to shape regularity. The combination of (67)–(69) yields

$$\begin{aligned} \|W_\star\|_{H^1(\Omega)}^2 &\lesssim \|(\mathbb{J}_\star - \mathbb{J}_\ell)g\|_{H^{1/2}(\Gamma_D)}^2 \lesssim \sum_{z \in \mathcal{K}_\ell^D \cap \omega_\ell^1} \|h_\ell^{1/2}(1 - \Pi_\ell) \nabla_\Gamma g\|_{L^2(\omega_{\ell,z}^4)}^2 \\ &\simeq \|h_\ell^{1/2}(1 - \Pi_\ell) \nabla_\Gamma g\|_{L^2(\omega_\ell^5)}^2 \\ &\lesssim \sum_{E \in \tilde{\mathcal{R}}_\ell^D} \text{osc}_{D,\ell}(E)^2, \end{aligned}$$

due to (41)–(42) in Proposition 8. Defining the set

$$\mathcal{R}_\ell := \mathcal{T}_\ell \setminus \mathcal{T}_\star \cup \{T \in \mathcal{T}_\ell : \exists E \in \tilde{\mathcal{R}}_\ell^D \quad E \subset \partial T\},$$

we observe $\mathcal{T}_\ell \setminus \mathcal{T}_\star \subseteq \mathcal{R}_\ell$ as well as $\#\mathcal{R}_\ell \lesssim \#(\mathcal{T}_\ell \setminus \mathcal{T}_\star)$. Moreover, the definition of the local contributions of $\tilde{\eta}_\ell$ in (50) shows

$$\sum_{E \in \tilde{\mathcal{R}}_\ell^D} \text{osc}_{D,\ell}(E)^2 \leq \sum_{T \in \mathcal{R}_\ell} \tilde{\eta}_\ell(T)^2.$$

This concludes the proof. \square

Corollary 22 (optimality of Dörfler marking for Scott-Zhang projection). *For arbitrary $0 < \kappa_\star < 1$, there is a constant $0 < \theta_\star < 1$ such that for all $\ell \in \mathbb{N}_0$ and all meshes $\mathcal{T}_\star = \text{refine}(\mathcal{T}_\ell)$ with $\tilde{\eta}_\star^2 \leq \kappa_\star \tilde{\eta}_\ell^2$, the set $\mathcal{R}_\ell \subseteq \mathcal{T}_\ell$ from Proposition 21 satisfies the Dörfler marking*

$$\theta \tilde{\eta}_\ell^2 \leq \sum_{T \in \mathcal{R}_\ell} \tilde{\eta}_\ell(T)^2 \quad (70)$$

for all $0 < \theta \leq \theta_\star$.

Proof. We split the estimator into the contributions on the non-refined resp. refined elements

$$\tilde{\eta}_\ell^2 = \sum_{T \in \mathcal{T}_\ell} \tilde{\eta}_\ell(T)^2 = \sum_{T \in \mathcal{T}_\ell \cap \mathcal{T}_\star} \tilde{\eta}_\ell(T)^2 + \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{T}_\star} \tilde{\eta}_\ell(T)^2.$$

Arguing as for the estimator reduction in Proposition 11 with $\delta = 1$, we see

$$\sum_{T \in \mathcal{T}_\ell \cap \mathcal{T}_\star} \tilde{\eta}_\ell(T)^2 \leq 2 \sum_{T \in \mathcal{T}_\ell \cap \mathcal{T}_\star} \tilde{\eta}_\star(T)^2 + C_{\text{red}} \|\nabla(\tilde{U}_\star - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 \leq 2\tilde{\eta}_\star^2 + C_{\text{red}} \|\nabla(\tilde{U}_\star - \tilde{U}_\ell)\|_{L^2(\Omega)}^2.$$

We now combine both estimates and use $\tilde{\eta}_\star^2 \leq \kappa_\star \tilde{\eta}_\ell^2$ as well as the discrete local reliability with $\mathcal{R}_\ell \supseteq \mathcal{T}_\ell \setminus \mathcal{T}_\star$ to see

$$\tilde{\eta}_\ell^2 \leq 2\tilde{\eta}_\star^2 + C_{\text{red}} \|\nabla(\tilde{U}_\star - \tilde{U}_\ell)\|_{L^2(\Omega)}^2 + \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{T}_\star} \tilde{\eta}_\ell(T)^2 \leq 2\kappa_\star \tilde{\eta}_\ell^2 + (C_{\text{red}}C_{\text{dlr}} + 1) \sum_{T \in \mathcal{R}_\ell} \tilde{\eta}_\ell(T)^2.$$

Rearranging this estimate, we obtain

$$(C_{\text{red}}C_{\text{dlr}} + 1)^{-1}(1 - 2\kappa_\star) \tilde{\eta}_\ell^2 \leq \sum_{T \in \mathcal{R}_\ell} \tilde{\eta}_\ell(T)^2,$$

so that $0 < \theta_\star := (C_{\text{red}}C_{\text{dlr}} + 1)^{-1}(1 - 2\kappa_\star) < 1$ concludes the proof. \square

6.2. Optimality of newest vertex bisection. The quasi-optimality analysis of AFEM requires two properties of the mesh-refinement which are satisfied for newest vertex bisection: First, for two triangulations $\mathcal{T}', \mathcal{T}'' \in \mathbb{T}$, let $\mathcal{T}' \oplus \mathcal{T}'' \in \mathbb{T}$ be the coarsest common refinement of both. Since newest vertex bisection is a binary refinement rule, it can be proved that $\mathcal{T}' \oplus \mathcal{T}''$ is just the overlay of both meshes, see [S07, Proof of Lemma 5.2] for 2D and the generalization to arbitrary dimension in [CKNS, Lemma 3.7]. Moreover, the number of elements of the overlay is controlled by

$$\#(\mathcal{T}' \oplus \mathcal{T}'') \leq \#\mathcal{T}' + \#\mathcal{T}'' - \#\mathcal{T}_0, \quad (71)$$

since both meshes are generated from the initial mesh \mathcal{T}_0 .

Second, we need the optimality of the mesh-closure, i.e. the definition $\mathcal{T}_{\ell+1} = \text{refine}(\mathcal{T}_\ell, \mathcal{M}_\ell)$ leads at least to refinement of all marked elements $T \in \mathcal{M}_\ell$. In addition, further elements

$T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell$ have to be refined to ensure conformity of the mesh. It has been proved in [BDD, Theorem 2.4] for 2D that

$$\#\mathcal{T}_{\ell+1} - \#\mathcal{T}_0 \leq C_{\text{nvb}} \sum_{j=0}^{\ell} \#\mathcal{M}_j \quad \text{for all } \ell \geq 0, \quad (72)$$

i.e. the number of elements in $\mathcal{T}_{\ell+1}$ is bounded by the number of marked elements. The constant $C_{\text{nvb}} > 0$ depends only on \mathcal{T}_0 in the sense that the initial reference edge distribution had to satisfy a certain assumption. Very recently [KPP], it could be proved that (72) holds without any further assumptions on \mathcal{T}_0 . For arbitrary dimension, (72) has been proved in [S08, Theorem 6.1] and \mathcal{T}_0 has to satisfy a certain assumption on the initial reference edge distribution.

6.3. Proof of quasi-optimality of AFEM (Theorem 6). In a first step, we prove that $\#\mathcal{M}_\ell \lesssim \Delta_\ell^{-1/(2s)}$. To that end, let $\varepsilon > 0$ be a free parameter which is determined later. According to the definition of the approximation class \mathbb{A}_s , there is some triangulation $\mathcal{T}_\varepsilon \in \mathbb{T}$ with

$$\eta_\varepsilon \leq \varepsilon \quad \text{and} \quad \#\mathcal{T}_\varepsilon - \#\mathcal{T}_0 \lesssim \varepsilon^{-1/s},$$

where the hidden constant depends only on \mathbb{A}_s . We consider the overlay $\mathcal{T}_\star := \mathcal{T}_\varepsilon \oplus \mathcal{T}_\ell$. Arguing as for the estimator reduction (52) and use of the discrete local reliability for $\tilde{\eta}_\ell$, we obtain

$$\tilde{\eta}_\star \lesssim \tilde{\eta}_\ell + \|\nabla(\tilde{U}_\star - \tilde{U}_\ell)\|_{L^2(\Omega)} \lesssim \tilde{\eta}_\ell \simeq \eta_\ell \leq \varepsilon,$$

where we have finally used the equivalence of both error estimators provided by Lemma 18. Choosing $\varepsilon = \delta \eta_\ell \simeq \delta \tilde{\eta}_\ell$ with sufficiently small $\delta > 0$, we thus infer

$$\tilde{\eta}_\star \leq \tilde{\kappa}_\star \tilde{\eta}_\ell$$

with some appropriate $0 < \tilde{\kappa}_\star \leq \kappa_\star$, where arbitrary $0 < \kappa_\star < 1$ in Proposition 22 fixes $0 < \theta_\star < 1$. The constant $\tilde{\kappa}_\star$ will be determined later. Together with the overlay estimate (71), we infer

$$\#\mathcal{R}_\ell \simeq \#(\mathcal{T}_\ell \setminus \mathcal{T}_\star) \leq \#\mathcal{T}_\star - \#\mathcal{T}_\ell \leq \#\mathcal{T}_\varepsilon - \#\mathcal{T}_0 \lesssim \varepsilon^{-1/s}$$

as well as the Dörfler estimate

$$\theta_\star \tilde{\eta}_\ell^2 \leq \sum_{T \in \mathcal{R}_\ell} \tilde{\eta}_\ell(T)^2.$$

We now need to show that this implies Stevenson's modified Dörfler marking. To that end, we again employ Lemma 18:

- In case of $\text{osc}_{D,\ell}^2 \leq \vartheta \varrho_\ell^2$, we employ Lemma 18 twice to see

$$\theta_\star \varrho_\ell^2 \lesssim \theta_\star \tilde{\eta}_\ell^2 \leq \sum_{T \in \mathcal{R}_\ell} \tilde{\eta}_\ell(T)^2 + \text{osc}_{D,\ell}^2 \lesssim \sum_{T \in \mathcal{R}_\ell} \varrho_\ell(T)^2 + \text{osc}_{D,\ell}^2 \leq \sum_{T \in \mathcal{R}_\ell} \varrho_\ell(T)^2 + \vartheta \varrho_\ell^2.$$

Put differently, we obtain

$$((C_{\text{eq}} + 1)^{-2} \theta_\star - \vartheta) \varrho_\ell^2 \leq \sum_{T \in \mathcal{R}_\ell} \varrho_\ell(T)^2,$$

i.e. for $0 < \vartheta, \theta_1 < 1$ sufficiently small, the set $\mathcal{R}_\ell \subseteq \mathcal{T}_\ell$ satisfies the marking criterion (11).

- In case of $\text{osc}_{D,\ell}^2 > \vartheta \varrho_\ell^2$, we use that the Dirichlet oscillations are locally determined, i.e.

$$\sum_{E \in \mathcal{E}_\ell^D \cap \mathcal{E}_\star^D} \text{osc}_{D,\ell}(E)^2 = \sum_{E \in \mathcal{E}_\ell^D \cap \mathcal{E}_\star^D} \text{osc}_{D,\star}(E)^2 \leq \text{osc}_{D,\star}^2 \leq \tilde{\eta}_\star^2 \leq \tilde{\kappa}_\star \tilde{\eta}_\ell^2 \simeq \tilde{\kappa}_\star \eta_\ell^2 \leq \tilde{\kappa}_\star (1 + \vartheta^{-1}) \text{osc}_{D,\ell}^2.$$

This estimate yields

$$(1 - \tilde{\kappa}_\star (1 + \vartheta^{-1})(C_{\text{eq}} + 1)) \text{osc}_{D,\ell}^2 \leq \sum_{E \in \mathcal{E}_\ell^D \setminus \mathcal{E}_\star^D} \text{osc}_{D,\ell}(E)^2$$

For arbitrary $0 < \theta_2 < 1$ and sufficiently small $0 < \tilde{\kappa}_\star < 1$, we infer that $\mathcal{E}_\ell^D \setminus \mathcal{E}_\star^D$ satisfies the marking criterion (12).

In the first case, minimal cardinality of $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell$ in step (iii) of Algorithm 1 implies $\#\mathcal{M}_\ell \leq \#\mathcal{R}_\ell \simeq \#(\mathcal{T}_\ell \setminus \mathcal{T}_\star)$. In the second case, minimal cardinality of $\mathcal{M}_\ell^D \subseteq \mathcal{E}_\ell^D$ and the definition of $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell$ in step (iv) of Algorithm 1 imply $\#\mathcal{M}_\ell \leq \#\mathcal{M}_\ell^D \leq \#(\mathcal{E}_\ell^D \setminus \mathcal{E}_\star^D) \lesssim \#(\mathcal{T}_\ell \setminus \mathcal{T}_\star)$. In either case, we thus conclude

$$\#\mathcal{M}_\ell \lesssim \#(\mathcal{T}_\ell \setminus \mathcal{T}_\star) \lesssim \varepsilon^{-1/s} \simeq \eta_\ell^{-1/s} \simeq \Delta_\ell^{-1/(2s)}. \quad \text{for all } \ell \geq 0.$$

We now conclude the proof as e.g. in [S07, CKNS]: By use of the closure estimate (72), we obtain

$$\#\mathcal{T}_\ell - \#\mathcal{T}_0 \lesssim \sum_{j=0}^{\ell-1} \#\mathcal{M}_j \lesssim \sum_{j=0}^{\ell-1} \Delta_j^{-1/(2s)}.$$

Note that the contraction property (20) of Δ_j implies $\Delta_\ell \leq \kappa^{\ell-j} \Delta_j$, whence $\Delta_j^{-1/(2s)} \leq \kappa^{(\ell-j)/(2s)} \Delta_\ell^{-1/(2s)}$. According to $0 < \kappa < 1$ and the geometric series, this gives

$$\#\mathcal{T}_\ell - \#\mathcal{T}_0 \lesssim \Delta_\ell^{-1/(2s)} \sum_{j=0}^{\ell-1} \kappa^{(\ell-j)/(2s)} \lesssim \Delta_\ell^{-1/(2s)} \simeq \eta_\ell^{-1/s}.$$

Altogether, we may therefore conclude that $(u, f, g, \phi) \in \mathbb{A}_s$ implies $\eta_\ell \lesssim (\#\mathcal{T}_\ell - \#\mathcal{T}_0)^{-s}$ for all $\ell \geq 0$. The converse implication is obvious by definition of \mathbb{A}_s . \square

6.4. Characterization of approximation class (Theorem 7). First, note that for a given mesh $\mathcal{T}_\star \in \mathbb{T}$ the estimator η_\star dominates all oscillation terms, i.e.

$$\text{osc}_{\mathcal{T},\star} \leq \eta_\star, \quad \text{osc}_{D,\star} \leq \eta_\star, \quad \text{osc}_{N,\star} \leq \eta_\star.$$

We assume $(u, f, g, \phi) \in \mathbb{A}_s$ for some $s > 0$. For each $N \in \mathbb{N}$ it exists $\mathcal{T}_\star \in \mathbb{T}_N$ such that

$$N^s \text{osc}_{\mathcal{T},\star} \leq N^s \eta_\star \leq C := \sup_{N \in \mathbb{N}} \inf_{\mathcal{T}_\star \in \mathbb{T}_N} N^s \eta_\star < \infty. \quad (73)$$

Analogously, we have

$$N^s \text{osc}_{N,\star} \leq C < \infty \quad \text{and} \quad N^s \text{osc}_{D,\star} \leq C < \infty. \quad (74)$$

The reliability result in Proposition 3 yields

$$\min_{V_\star \in \mathcal{S}^p(\mathcal{T}_\star)} N^s \|u - V_\star\|_{H^1(\Omega)} \leq N^s \|u - U_\star\|_{H^1(\Omega)} \leq C_{\text{rel}} N^s \eta_\star \leq C_{\text{rel}} C < \infty. \quad (75)$$

Because $N \in \mathbb{N}$ was arbitrary, the estimates (73)–(75) prove (26)–(29).

Now, we assume that (26)–(29) hold for (u, f, g, ϕ) . We aim to prove $(u, f, g, \phi) \in \mathbb{A}_s$. By use of the efficiency estimate in Proposition 3 and the Céa-type estimate in Proposition 2, we derive

$$\begin{aligned} \sup_{N \in \mathbb{N}} \inf_{\mathcal{T}_* \in \mathbb{T}_N} N^s \eta_* &\leq C_{\text{eff}} \sup_{N \in \mathbb{N}} \inf_{\mathcal{T}_* \in \mathbb{T}_N} N^s \left(C_{\text{Céa}} \min_{V_* \in \mathcal{S}^p(\mathcal{T}_*)} \|u - V_*\|_{H^1(\Omega)}^2 \right. \\ &\quad \left. + \text{osc}_{\mathcal{T},*}^2 + \text{osc}_{N,*}^2 + \text{osc}_{D,*}^2 \right). \end{aligned} \quad (76)$$

For $N \in \mathbb{N}$, the assumption (26)–(29) guarantee meshes $\mathcal{T}_{*u}, \mathcal{T}_{*\Omega}, \mathcal{T}_{*N}, \mathcal{T}_{*D} \in \mathbb{T}_{N/4}$ such that

$$\begin{aligned} (N/4)^s \min_{V_{*u} \in \mathcal{S}^p(\mathcal{T}_{*u})} \|u - V_{*u}\|_{H^1(\Omega)} &\leq \sup_{N > 0} \inf_{\mathcal{T}_* \in \mathbb{T}_N} \min_{V_* \in \mathcal{S}^p(\mathcal{T}_*)} N^s \|u - V_*\|_{H^1(\Omega)} =: C_u < \infty, \\ (N/4)^s \text{osc}_{\mathcal{T},*\Omega} &\leq \sup_{N > 0} \inf_{\mathcal{T}_* \in \mathbb{T}_N} N^s \text{osc}_{\mathcal{T},*} =: C_{\text{osc}_{\mathcal{T}}} < \infty, \\ (N/4)^s \text{osc}_{N,*N} &\leq \sup_{N > 0} \inf_{\mathcal{T}_* \in \mathbb{T}_N} N^s \text{osc}_{N,*} =: C_{\text{osc}_N} < \infty, \\ (N/4)^s \text{osc}_{D,*D} &\leq \sup_{N > 0} \inf_{\mathcal{T}_* \in \mathbb{T}_N} N^s \text{osc}_{D,*} =: C_{\text{osc}_D} < \infty. \end{aligned}$$

Now, we consider the overlay $\mathcal{T}_* := \mathcal{T}_{*u} \oplus \mathcal{T}_{*\Omega} \oplus \mathcal{T}_{*N} \oplus \mathcal{T}_{*D}$. The overlay estimate (71) gives $\#\mathcal{T}_* \leq N - 3\#\mathcal{T}_0$, whence $\#\mathcal{T}_* - \#\mathcal{T}_0 \leq N$. Due to the fact that Π_E and Π_T are projections, we get immediately by definition of the oscillation terms and $\mathcal{S}^p(\mathcal{T}_*) \supseteq \mathcal{S}^p(\mathcal{T}_{*u})$

$$\begin{aligned} \min_{V_* \in \mathcal{S}^p(\mathcal{T}_*)} \|u - V_*\|_{H^1(\Omega)} &\leq \min_{V_{*u} \in \mathcal{S}^p(\mathcal{T}_{*u})} \|u - V_{*u}\|_{H^1(\Omega)}, \\ \text{osc}_{\mathcal{T},*} &\leq \text{osc}_{\mathcal{T},*\Omega}, \quad \text{osc}_{N,*} \leq \text{osc}_{N,*N}, \quad \text{and} \quad \text{osc}_{D,*} \leq \text{osc}_{D,*D}. \end{aligned}$$

Together with (76), we prove

$$\begin{aligned} \inf_{\mathcal{T}_* \in \mathbb{T}_N} N^s \eta_* &\leq C_{\text{eff}} N^s \left(C_{\text{Céa}} \min_{V_* \in \mathcal{S}^p(\mathcal{T}_*)} \|u - V_*\|_{H^1(\Omega)}^2 + \text{osc}_{\mathcal{T},*}^2 + \text{osc}_{N,*}^2 + \text{osc}_{D,*}^2 \right) \\ &\leq C_{\text{eff}} N^s \left(C_{\text{Céa}} \min_{V_{*u} \in \mathcal{S}^p(\mathcal{T}_{*u})} \|u - V_{*u}\|_{H^1(\Omega)}^2 + \text{osc}_{\mathcal{T},*\Omega}^2 + \text{osc}_{N,*N}^2 + \text{osc}_{D,*D}^2 \right) \\ &\leq C_{\text{eff}} 4^{-s} (C_{\text{Céa}} C_u + C_{\text{osc}_{\mathcal{T}}} + C_{\text{osc}_N} + C_{\text{osc}_D}) < \infty, \end{aligned}$$

where the constants are independent of $N \in \mathbb{N}$. Taking the supremum over $N \in \mathbb{N}$, we conclude $(u, f, g, \phi) \in \mathbb{A}_s$. \square

7. NUMERICAL EXPERIMENT


In this section, we provide numerical results for mixed boundary value problems in two and three space dimensions for the lowest-order case $p = 1$. In both examples we choose $\vartheta = \theta_1 = \theta_2$ in Algorithm 1. For comparison of the individual contributions ϱ_ℓ , we further define the jump terms

$$\varrho_\ell(\mathcal{E}_\ell^\Omega)^2 := \sum_{E \in \mathcal{E}_\ell^\Omega} |T|^{1/d} \|[\partial_n U_\ell]\|_{L^2(E)}^2, \quad (77)$$

the volume terms

$$\varrho_\ell(\Omega)^2 := \sum_{T \in \mathcal{T}_\ell} |T|^{2/d} \|f\|_{L^2(T)}^2 \quad (78)$$

Error estimator ϱ_ℓ for L^2 -orthogonal projection in 2D



Error estimator ϱ_ℓ for Scott-Zhang projection in 2D




FIGURE 1. Numerical results for ϱ_ℓ for uniform and adaptive mesh-refinement using $\theta \in \{1/4, 1/8, 1/16\}$ and L^2 -orthogonal projection and Scott-Zhang projection in 2D, respectively, plotted over the number of elements $N = \#\mathcal{T}_\ell$.

Individual contributions of ϱ_ℓ for Scott-Zhang projection in 2D




FIGURE 2. Numerical results for $\varrho_\ell(\mathcal{E}_\ell^\Omega)$, $\varrho_\ell(\mathcal{E}_\ell^N)$ and $\text{osc}_{D,\ell}$ for uniform and adaptive mesh-refinement using $\theta = 0.25$ and Scott-Zhang projection in 2D, plotted over the number of elements $N = \#\mathcal{T}_\ell$.

and the Neumann terms

$$\varrho_\ell(\mathcal{E}_\ell^N)^2 := \sum_{E \in \mathcal{E}_\ell^N} |T|^{1/d} \|\phi - \partial_n U_\ell\|_{L^2(E)}^2. \quad (79)$$

for the respective space dimension $d \in 2, 3$.

7.1. 2D example on Z-shape. In our first example, we consider the Z-shaped domain $\Omega = (-1, 1)^2 \setminus \text{conv}\{(0, 0), (-1, -1), (0, -1)\}$, see Figure 3, where also the partition of the boundary $\Gamma = \partial\Omega$ into Dirichlet boundary Γ_D and Neumann boundary Γ_N as well as the initial mesh is shown. We prescribe the exact solution

$$u(x) = r^{4/7} \cos(4\varphi/7) \quad (80)$$

of problem 1 in polar coordinates $x = r(\cos \varphi, \sin \varphi)$ and compute the Neumann and Dirichlet data thereof. Note, that f is harmonic so that

$$-\Delta u = f = 0.$$

The solution u as well as the Dirichlet data $g = u|_\Gamma$ show a generic singularity at the reentrant corner $r = 0$.




FIGURE 3. Z-shaped domain with initial mesh \mathcal{T}_0 and adaptively generated mesh \mathcal{T}_{11} with $N = 9.864$ for $\theta = 0.25$. The Dirichlet boundary Γ_D is marked red, whereas the blue parts denote the Neumann boundary $\Gamma \setminus \Gamma_D$.

Figure 1 shows a comparison between uniform and adaptive mesh refinement, where the adaptivity parameter θ varies between $1/4$ and $1/16$ and where the Dirichlet data are discretized by means of the L^2 -projection and the Scott-Zhang projection, respectively. It is easily seen that both discretizations lead to the optimal convergence rate $\mathcal{O}(N^{-1/2})$ for all parameters θ , whereas uniform refinement leads only to suboptimal convergence behaviour of approximately $\mathcal{O}(N^{-2/7})$. Note that due to $f \equiv 0$, we have no volume contributions in this example.

In Figure 2, we compare the jump terms, the Neumann terms, as well as the Dirichlet oscillations $\text{osc}_{D,\ell}$ for uniform and adaptive refinement, where we have chosen the Scott-Zhang projection to discretize the boundary data. Even here, we observe better convergence rates with adaptive refinement. Due to the corner singularity of the exact solution at $r = 0$, uniform refinement leads to a suboptimal convergence behaviour, even for the oscillations. Finally, in Figure 3, the initial mesh \mathcal{T}_0 and the adaptively generated mesh \mathcal{T}_{11} with $N = 9.864$ Elements are visualized. As expected, adaptive refinement is essentially concentrated around the reentrant corner $r = 0$.

7.2. 3D example on the Fichera cube. As computational domain serves the Fichera cube $\Omega = (-1, 1)^3 \setminus [0, 1]^3$ which has a concave corner and three reentrant edges. The partition of the boundary $\Gamma = \partial\Omega$ into Dirichlet boundary Γ_D and Neumann boundary Γ_N , as well as the initial surface mesh is shown in Figure 6. We solve problem (1) with right-hand side

$$f(x, y, z) := -\frac{5}{16} (x^2 + y^2 + z^2)^{-7/8}.$$

The boundary data are prescribed by the trace resp. normal derivative of the exact solution

$$u(x, y, z) = (x^2 + y^2 + z^2)^{1/8}$$





FIGURE 4. Numerical results for ϱ_ℓ for uniform and adaptive mesh-refinement using $\theta \in \{1/4, 1/8, 1/16\}$ and L^2 -orthogonal projection and Scott-Zhang projection in 3D, respectively, plotted over the number of elements $N = \#\mathcal{T}_\ell$.

Individual contributions of ϱ_ℓ for L^2 -orthogonal projection in 3D




FIGURE 5. Numerical results for $\varrho_\ell(\Omega)$, $\varrho_\ell(\mathcal{E}_\ell^\Omega)$, $\varrho_\ell(\mathcal{E}_\ell^N)$ and $\text{osc}_{D,\ell}$ for uniform and adaptive mesh-refinement using $\theta = 1/4$ and L^2 -orthogonal projection in 3D, plotted over the number of elements $N = \#\mathcal{T}_\ell$.

which has a singular gradient at the reentrant corner at the origin. Similar to the 2D case, we provide comparisons for various adaptivity parameters as well as for different choices for the discretization of the boundary data.

In Figure 4, we compare uniform and adaptive mesh refinement where the Dirichlet data are discretized by means of the L^2 -orthogonal projection or the Scott-Zhang projection, respectively. The adaptivity parameter is varied between $1/4$ and $1/16$. We observe that either discretization g_ℓ of the Dirichlet data g leads to the optimal convergence rate $\mathcal{O}(N^{-1/3})$ for all choices of θ . Due to the generic singularity at the center, uniform refinement leads only to suboptimal convergence rate of $\mathcal{O}(N^{-2/9})$.

In Figure 5, we compare each contribution of the estimator separately for uniform and adaptive refinement with $\theta = 1/4$. For this comparison, we chose the L^2 -orthogonal projection to discretize g . For adaptive refinement, we observe optimal order of convergence even for $\varrho_\ell(\mathcal{E}_\ell^\Omega)$, $\varrho_\ell(\Omega)$, $\varrho_\ell(\mathcal{E}_\ell^N)$, and $\text{osc}_{D,\ell}$. Uniform refinement, on the other hand, leads to suboptimal convergence rate also for the individual contributions.




FIGURE 6. Fichera cube with boundary of the initial mesh \mathcal{T}_0 and \mathcal{T}_{30} with $N = 200.814$ for $\theta = 0.25$. The Dirichlet boundary $\Gamma_D = \{-1\} \times [-1, 1]^2$ is marked red, whereas the blue parts denote the Neumann boundary $\Gamma \setminus \Gamma_D$.

The computational domain, with initial (surface) mesh \mathcal{T}_0 as well as the adaptively generated mesh \mathcal{T}_{30} with $\#\mathcal{T}_{30} = 200.814$ elements is finally shown in Figure 6. As expected, the refinement is basically concentrated around the singularity at the origin.

Acknowledgements The authors M.A., D.P., and M.F. are funded by the Austrian Science Fund (FWF) through grant P21732 “*Adaptive Boundary Element Method*”. The author M.P. acknowledges support from the Viennese Science and Technology Fund (WWTF) under grant MA09-029 “*Micromagnetic Simulation and Computational Design of Future Devices*”.

REFERENCES

- [AFP] M. AURADA, S. FERRAZ-LEITE, D. PRAETORIUS: *Estimator reduction and convergence of adaptive BEM*, Appl. Numer. Math., in press (2011).
- [AO] M. AINSWORTH, T. ODEN: *A posteriori error estimation in finite element analysis*, Wiley-Interscience, New-York, 2000.
- [BCD] S. BARTELS, C. CARSTENSEN, G. DOLZMANN: *Inhomogeneous Dirichlet conditions in a priori and a posteriori finite element error analysis*, Numer. Math. **99** (2004), 1–24.
- [BDD] P. BINEV, W. DAHMEN, R. DEVORE: *Adaptive finite element methods with convergence rates*, Numer. Math. **97** (2004), 219–268.
- [BM] R. BECKER, S. MAO: *Convergence and quasi-optimal complexity of a simple adaptive finite element method*, M2AN Math. Model. Numer. Anal. **43** (2009), 1203–1219.
- [BV] I. BABUŠKA, M. VOGELIUS: *Feedback and adaptive finite element solution of one-dimensional boundary value problems*, Numer. Math. **44** (1984), 75–102.
- [CMS] C. CARSTENSEN, M. MAISCHAK, E.P. STEPHAN: *A posteriori error estimate and h-adaptive algorithm on surfaces for Symm’s integral equation*, Numer. Math. **90** (2001), no. 2, 197–213.
- [CKNS] M. CASCÓN, C. KREUZER, R. NOCHETTO, K. SIEBERT: *Quasi-optimal convergence rate for an adaptive finite element method*, SIAM J. Numer. Anal. **46** (2008), 2524–2550.
- [D] W. DÖRFLER: *A convergent adaptive algorithm for Poisson’s equation*, SIAM J. Numer. Anal. **33** (1996), 1106–1124.

- [FKMP] M. FEISCHL, M. KARKULIK, M. MELENK, D. PRAETORIUS: *Quasi-optimal convergence rate for an adaptive boundary element method*, ASC Report **28/2011**, Institute for Analysis and Scientific Computing, Vienna University of Technology, 2011.
- [FPP] M. FEISCHL, M. PAGE, D. PRAETORIUS: *Convergence and quasi-optimality of adaptive FEM with inhomogeneous Dirichlet data*, ASC Report **34/2010**, Institute for Analysis and Scientific Computing, Vienna University of Technology, 2010.
- [KS] C. KREUZER, K. SIEBERT: *Decay rates of adaptive finite elements with Dörfler marking*, Numer. Math. **117** no. 4 (2011), 679–716.
- [KOP] M. KARKULIK, G. OF, D. PRAETORIUS: *Convergence of adaptive 3D BEM for some weakly-singular integral equation*, work in progress (2011).
- [KPP] M. KARKULIK, D. PAVLICEK, D. PRAETORIUS: *On newest vertex bisection in 2D*, work in progress (2011).
- [MNS] P. MORIN, R. NOCHETTO, K. SIEBERT: *Data oscillation and convergence of adaptive FEM*, SIAM J. Numer. Anal. **18** (2000), 466–488.
- [MSV] P. MORIN, K. SIEBERT, A. VEESER: *A basic convergence result for conforming adaptive finite elements*, Math. Models Methods Appl. Sci. **18** (2008), 707–737.
- [SV] R. SACCHI, A. VEESER: *Locally efficient and reliable a posteriori error estimators for Dirichlet problems*, Math. Models Methods Appl. Sci. **16** (2006), 319–346.
- [SZ] L. SCOTT, S. ZHANG: *Finite element interpolation of nonsmooth functions satisfying boundary conditions*, Math. Comp. **54** (1990), 483–493.
- [S07] R. STEVENSON: *Optimality of standard adaptive finite element method*, Found. Comput. Math. (2007), 245–269.
- [S08] R. STEVENSON: *The completion of locally refined simplicial partitions created by bisection*, Math. Comp. **77** (2008), 227–241.
- [T] TRAXLER: *An Algorithm for Adaptive Mesh Refinement in n Dimensions*, Computing **59** (1997), 115–137.
- [V] R. VERFÜRTH: *A review of a posteriori error estimation and adaptive mesh-refinement techniques*, Wiley-Teubner, 1996.

INSTITUTE FOR ANALYSIS AND SCIENTIFIC COMPUTING, VIENNA UNIVERSITY OF TECHNOLOGY, WIEDNER HAUPTSTRASSE 8-10, A-1040 WIEN, AUSTRIA

E-mail address: {Markus.Aurada, Michael.Feischl, Josef.Kemetmueller, Dirk.Praetorius}@tuwien.ac.at

E-mail address: Marcus.Page@tuwien.ac.at (corresponding author)